VOLUME 38 • AUTHOR INDEX

Aaltonen LA *see* Aaltonen L-M *et al*
see Avizienyte E *et al*
Aaltonen L-M *et al*. Role of *TP*53 P72R polymorphism in human papillomavirus associated premalignant laryngeal neoplasm: *letter*, **327**
Abel A *see* Chompret A *et al*
Adams G *see* Talks SJ *et al*
Adèr HJ *see* Henneman L *et al*
Adoun M *see* Amaral MD *et al*
Affara NA *see* Boucher CA *et al*
see Ross NLJ *et al*
Ahmed M *see* Henwood J *et al*
Ahmed Z *see* Ben-Yosef T *et al*
Ahmed ZM *see* Ben-Yosef T *et al*
Ahonen T *see* Nopola-Hemmi J *et al*
Aittomäki K *see* Gaff C *et al*
Akaboshi S *see* Yamamoto T *et al*
Akbas F *et al*. Molecular and clinical study of two myotonic dystrophy homozygotes: *letter*, **e40**
Akrami SM *et al*. Detection of a large *TBX5* deletion in a family with Holt-Oram syndrome, **e44**
Aktan-Collan K *et al*. Comprehension of cancer risk one and 12 months after predictive genetic testing for hereditary non-polyposis colorectal cancer: *letter*, **787**
Albers A *see* Eggermann T *et al*
Albrecht B *et al*. Uniparental isodisomy for paternal 2p and maternal 2q in a phenotypically normal female with two isochromosomes, i(2p) and i(2q): *letter*, **214**
Al-Din A *see* Hampshire DJ *et al*
Al-Gazali LI *see* Bayoumi R *et al*
Allada V *see* Vitale E *et al*
Alonso LG *see* Nadal M *et al*
Altehöfer C *see* Bender BU *et al*
Alvarado D *see* Rainier S *et al*
Amaral MD *et al*. Cystic fibrosis patients with the 3272-26A>G splicing mutation have milder disease than F508del homozygotes: a large European study, **777**
Ameen M *see* Veal CD *et al*
Amidon D *see* Lossie AC *et al*
Amiel J *see* Imessaoudene B *et al*
Amiel J, Lyonnet S. Hirschsprung disease, associated syndromes, and genetics: a review: *review*, **729**
Amiel J *et al*. Trinucleotide repeat contraction: a pitfall in prenatal diagnosis of myotonic dystrophy : *letter*, **850**
Andersen PS *et al*. Myosin light chain mutations in familial hypertrophic cardiomyopathy: phenotypic presentation and frequency in Danish and South African populations, **e43**
Andersen PS *see* Mogensen J *et al*
Andrade JAD *see* Nadal M *et al*
Angelopoulou R *see* Siffroi JP *et al*
Annuss G *see* Tönnies H *et al*
Antiñolo G *see* Li Y *et al*
Antoccia A *see* Maraschio P *et al*
Antonarakis SE *see* Ben-Yosef T *et al*
Antoniuk SA *see* Kwok JBJ *et al*
Anvret M *see* Villard L *et al*
Arias IM *see* Kadakol A *et al*
Aricó M *see* Clementi R *et al*
Ariga T *see* Kobayashi I *et al*
Armour JAL *see* Akrami SM *et al*
Arpaia E *see* Grunebaum E *et al*
Arte S *see* Pirinen S *et al*
Arveiler B *see* Coupry I *et al*
Aslan M *see* Fritz B *et al*
Au K-S *see* Khare L *et al*
Aubert D *see* Giraudeau F *et al*
Aubry MC *see* Viot G *et al*
Aurias A *see* Gad S *et al*
Ausems CWST *see* Ausems MGEM *et al*
Ausems MGEM *see* van der Luijt RB *et al*
Ausems MGEM *et al*. Dutch patients with glycogen storage disease type II show common ancestry for the 525delT and del exon 18 mutations: *letter*, **527**
Autere J *see* Finnilä S *et al*
Avizienyte E *et al*. *E-cadherin* is not frequently mutated in hereditary gastric cancer: *letter*, **49**
Aysal F *see* Akbas F *et al*
Ayuso C *see* Lorda-Sanchez I *et al*

Badiali M *see* Steigerwald K *et al*
Bahuau M *et al*. *GDNF* as a candidate modifier in a type 1 neurofibromatosis (NF1) enteric phenotype: *letter*, **638**
Bailey IC *see* McConnell RS *et al*
Bailey JN *see* Khare L *et al*
Baker L *et al*. Chromosome 2 interstitial deletion (del(2)(q14.1q21)) associated with connective tissue laxity and an attention deficit disorder: *letter*, **493**
Baldissera I *see* Janecke AR *et al*
Balendran N *see* Veal CD *et al*
Balko MG *see* Steigerwald K *et al*
Ballabio A *see* Kayserili H *et al*
Balmer D *see* Röthlisberger B *et al*
Barber JCK *see* Sumption ND and Barber JCK
Barber RC *see* Veal CD *et al*
Barberá R *see* Moreno-Pelayo MA *et al*
Bardoel AFJ *see* Ausems MGEM *et al*
Barkardottir RB *see* Bergthorsson JT *et al*
Barker JNWN *see* Veal CD *et al*
Barker K *et al*. *PTEN* mutations are uncommon in Proteus syndrome, **480**
Barlow-Stewart K *see* Meiser B *et al*
Barreto C *see* Amaral MD *et al*
Barros F *see* Ruiz-Ponte C *et al*
Barrot S *see* de Lonlay P *et al*
Barrow M *see* Watson P *et al*
Barton DE *see* Rowland JS *et al*
Barton J *see* Tobias S *et al*
Basaran S *see* Kayserili H *et al*
Baser ME *et al*. Maternal gene effect in neurofibromatosis 2: fact or artefact?: *letter*, **783**
Bateman A *see* Eccles D *et al*
Bates AW. A case of Roberts syndrome described in 1737: *letter*, **565**
Batista A *see* Seixas S *et al*
Baty CJ *see* Blair E *et al*
Baumann C *see* Faivre L *et al*
Bayoumi R *et al*. Localisation of a gene for an autosomal recessive syndrome of macrocephaly, multiple epiphyseal dysplasia, and distinctive facies to chromosome 15q26, **369**
Beals RK, LaFranchi S. Distal arthrogryposis: a new type with distinct facial appearance and absent teeth: *letter*, **e22**
Beck S *see* Amaral MD *et al*
Bedeschi M *see* Rossi E *et al*
Begnini A *see* Castellani C *et al*
Beis MJ *see* Coulter-Mackie MB *et al*
Bekiesiń-Figatowska MB *see* Chrzanowska KH *et al*
Benda TA *see* Mahavni V *et al*
Bender BU *et al*. *VHL* c.505 T>C mutation confers a high age related penetrance but no increased overall mortality, **508**
Benetazzo MG *see* Castellani C *et al*
Bennett CP *see* Henwood J *et al*
Bensimon A *see* Gad S *et al*
Ben-Yishay M *see* Dar P *et al*
Ben-Yosef T *et al*. Novel mutations of *TMPRSS3* in four DFNB8/B10 families segregating congenital autosomal recessive deafness: *letter*, **396**
Benzacken B *see* Siffroi JP *et al*
Berger DP *see* Bender BU *et al*
Bergthorsson JT *et al*. *BRCA1* and *BRCA2* mutation status and cancer family history of Danish women affected with multifocal or bilateral breast cancer at a young age, **361**
Berry PJ *see* Kelly AL *et al*
Berthaut I *see* Siffroi JP *et al*
Besana D *see* Maraschio P *et al*
Besmond C *see* Gabolde M *et al*
Bevan S *see* Barker K *et al*
Bhattacharya SS *see* Payne AM *et al*
Białecka M *see* Chrzanowska KH *et al*
Biancalana V *see* Giraudeau F *et al*
Bienvenu T *see* Amaral MD *et al*
see Bourdon V *et al*
Bierla J *see* Jaruzelska J *et al*
Bik E *see* Wagner A *et al*
Billingsley G *see* Vincent A *et al*
Bird AC *see* Payne AM *et al*
see Tarttelin EE *et al*
Bird L *see* Vitale E *et al*
Birembaut P *see* Amaral MD *et al*
Biswas P *see* Dollfus H *et al*
Bitner-Glindzicz M *see* Houseman MJ *et al*
Bitner-Glindzicz M *see* Rickard S *et al*
see Hutchin TP *et al*
see Kelsell DP *et al*
Black G *see* Watson P *et al*
Black S *see* Ungaro P *et al*
Blackburn J *see* Tarttelin EE *et al*
Blackman V *see* Ehrlich M *et al*
Blair E *et al*. Mutations in *cis* can confound genotype-phenotype correlations in hypertrophic cardiomyopathy: *letter*, **385**

VOLUME 38 • AUTHOR INDEX

Aaltonen LA *see* Aaltonen L-M *et al*
see Avizienyte E *et al*
Aaltonen L-M *et al*. Role of *TP*53 P72R polymorphism in human papillomavirus associated premalignant laryngeal neoplasm: *letter*, **327**
Abel A *see* Chompret A *et al*
Adams G *see* Talks SJ *et al*
Adèr HJ *see* Henneman L *et al*
Adoun M *see* Amaral MD *et al*
Affara NA *see* Boucher CA *et al*
see Ross NLJ *et al*
Ahmed M *see* Henwood J *et al*
Ahmed Z *see* Ben-Yosef T *et al*
Ahmed ZM *see* Ben-Yosef T *et al*
Ahonen T *see* Nopola-Hemmi J *et al*
Aittomäki K *see* Gaff C *et al*
Akaboshi S *see* Yamamoto T *et al*
Akbas F *et al*. Molecular and clinical study of two myotonic dystrophy homozygotes: *letter*, **e40**
Akrami SM *et al*. Detection of a large *TBX5* deletion in a family with Holt-Oram syndrome, **e44**
Aktan-Collan K *et al*. Comprehension of cancer risk one and 12 months after predictive genetic testing for hereditary non-polyposis colorectal cancer: *letter*, **787**
Albers A *see* Eggermann T *et al*
Albrecht B *et al*. Uniparental isodisomy for paternal 2p and maternal 2q in a phenotypically normal female with two isochromosomes, i(2p) and i(2q): *letter*, **214**
Al-Din A *see* Hampshire DJ *et al*
Al-Gazali LI *see* Bayoumi R *et al*
Allada V *see* Vitale E *et al*
Alonso LG *see* Nadal M *et al*
Altehöfer C *see* Bender BU *et al*
Alvarado D *see* Rainier S *et al*
Amaral MD *et al*. Cystic fibrosis patients with the 3272-26A>G splicing mutation have milder disease than F508del homozygotes: a large European study, **777**
Ameen M *see* Veal CD *et al*
Amidon D *see* Lossie AC *et al*
Amiel J *see* Imessaoudene B *et al*
Amiel J, Lyonnet S. Hirschsprung disease, associated syndromes, and genetics: a review: *review*, **729**
Amiel J *et al*. Trinucleotide repeat contraction: a pitfall in prenatal diagnosis of myotonic dystrophy : *letter*, **850**
Andersen PS *et al*. Myosin light chain mutations in familial hypertrophic cardiomyopathy: phenotypic presentation and frequency in Danish and South African populations, **e43**
Andersen PS *see* Mogensen J *et al*
Andrade JAD *see* Nadal M *et al*
Angelopoulou R *see* Siffroi JP *et al*
Annuss G *see* Tönnies H *et al*
Antiñolo G *see* Li Y *et al*
Antoccia A *see* Maraschio P *et al*
Antonarakis SE *see* Ben-Yosef T *et al*
Antoniuk SA *see* Kwok JBJ *et al*
Anvret M *see* Villard L *et al*
Arias IM *see* Kadakol A *et al*
Aricó M *see* Clementi R *et al*
Ariga T *see* Kobayashi I *et al*
Armour JAL *see* Akrami SM *et al*
Arpaia E *see* Grunebaum E *et al*
Arte S *see* Pirinen S *et al*
Arveiler B *see* Coupry I *et al*
Aslan M *see* Fritz B *et al*
Au K-S *see* Khare L *et al*
Aubert D *see* Giraudeau F *et al*
Aubry MC *see* Viot G *et al*
Aurias A *see* Gad S *et al*
Ausems CWST *see* Ausems MGEM *et al*
Ausems MGEM *see* van der Luijt RB *et al*
Ausems MGEM *et al*. Dutch patients with glycogen storage disease type II show common ancestry for the 525delT and del exon 18 mutations: *letter*, **527**
Autere J *see* Finnilä S *et al*
Avizienyte E *et al*. *E-cadherin* is not frequently mutated in hereditary gastric cancer: *letter*, **49**
Aysal F *see* Akbas F *et al*
Ayuso C *see* Lorda-Sanchez I *et al*

Badiali M *see* Steigerwald K *et al*
Bahuau M *et al*. *GDNF* as a candidate modifier in a type 1 neurofibromatosis (NF1) enteric phenotype: *letter*, **638**
Bailey IC *see* McConnell RS *et al*
Bailey JN *see* Khare L *et al*
Baker L *et al*. Chromosome 2 interstitial deletion (del(2)(q14.1q21)) associated with connective tissue laxity and an attention deficit disorder: *letter*, **493**
Baldissera I *see* Janecke AR *et al*
Balendran N *see* Veal CD *et al*
Balko MG *see* Steigerwald K *et al*
Ballabio A *see* Kayserili H *et al*
Balmer D *see* Röthlisberger B *et al*
Barber JCK *see* Sumption ND and Barber JCK
Barber RC *see* Veal CD *et al*
Barberá R *see* Moreno-Pelayo MA *et al*
Bardoel AFJ *see* Ausems MGEM *et al*
Barkardottir RB *see* Bergthorsson JT *et al*
Barker JNWN *see* Veal CD *et al*
Barker K *et al*. *PTEN* mutations are uncommon in Proteus syndrome, **480**
Barlow-Stewart K *see* Meiser B *et al*
Barreto C *see* Amaral MD *et al*
Barros F *see* Ruiz-Ponte C *et al*
Barrot S *see* de Lonlay P *et al*
Barrow M *see* Watson P *et al*
Barton DE *see* Rowland JS *et al*
Barton J *see* Tobias S *et al*
Basaran S *see* Kayserili H *et al*
Baser ME *et al*. Maternal gene effect in neurofibromatosis 2: fact or artefact?: *letter*, **783**
Bateman A *see* Eccles D *et al*
Bates AW. A case of Roberts syndrome described in 1737: *letter*, **565**
Batista A *see* Seixas S *et al*
Baty CJ *see* Blair E *et al*
Baumann C *see* Faivre L *et al*
Bayoumi R *et al*. Localisation of a gene for an autosomal recessive syndrome of macrocephaly, multiple epiphyseal dysplasia, and distinctive facies to chromosome 15q26, **369**
Beals RK, LaFranchi S. Distal arthrogryposis: a new type with distinct facial appearance and absent teeth: *letter*, **e22**
Beck S *see* Amaral MD *et al*
Bedeschi M *see* Rossi E *et al*
Begnini A *see* Castellani C *et al*
Beis MJ *see* Coulter-Mackie MB *et al*
Bekiesiń-Figatowska MB *see* Chrzanowska KH *et al*
Benda TA *see* Mahavni V *et al*
Bender BU *et al*. *VHL* c.505 T>C mutation confers a high age related penetrance but no increased overall mortality, **508**
Benetazzo MG *see* Castellani C *et al*
Bennett CP *see* Henwood J *et al*
Bensimon A *see* Gad S *et al*
Ben-Yishay M *see* Dar P *et al*
Ben-Yosef T *et al*. Novel mutations of *TMPRSS3* in four DFNB8/B10 families segregating congenital autosomal recessive deafness: *letter*, **396**
Benzacken B *see* Siffroi JP *et al*
Berger DP *see* Bender BU *et al*
Bergthorsson JT *et al*. *BRCA1* and *BRCA2* mutation status and cancer family history of Danish women affected with multifocal or bilateral breast cancer at a young age, **361**
Berry PJ *see* Kelly AL *et al*
Berthaut I *see* Siffroi JP *et al*
Besana D *see* Maraschio P *et al*
Besmond C *see* Gabolde M *et al*
Bevan S *see* Barker K *et al*
Bhattacharya SS *see* Payne AM *et al*
Białecka M *see* Chrzanowska KH *et al*
Biancalana V *see* Giraudeau F *et al*
Bienvenu T *see* Amaral MD *et al*
see Bourdon V *et al*
Bierla J *see* Jaruzelska J *et al*
Bik E *see* Wagner A *et al*
Billingsley G *see* Vincent A *et al*
Bird AC *see* Payne AM *et al*
see Tarttelin EE *et al*
Bird L *see* Vitale E *et al*
Birembaut P *see* Amaral MD *et al*
Biswas P *see* Dollfus H *et al*
Bitner-Glindzicz M *see* Houseman MJ *et al*
Bitner-Glindzicz M *see* Rickard S *et al*
see Hutchin TP *et al*
see Kelsell DP *et al*
Black G *see* Watson P *et al*
Black S *see* Ungaro P *et al*
Blackburn J *see* Tarttelin EE *et al*
Blackman V *see* Ehrlich M *et al*
Blair E *et al*. Mutations in *cis* can confound genotype-phenotype correlations in hypertrophic cardiomyopathy: *letter*, **385**

Blakley P *et al.* A clinical study of patients with multiple isolated neurofibromas: *letter*, **485**
Blancato J *see* Stratakis CA *et al*
Blanchet P *see* Lefort G *et al*
Blayau M *see* Amaral MD *et al*
Blescia A *see* Gong W *et al*
Boardman JP *et al.* A novel mutation in the endothelin B receptor gene in a patient with Shah-Waardenburg syndrome and Down syndrome: *letter*, **646**
Bobrow M *see* Michie S *et al*
Boer H *see* Whittington JE *et al*
Boetes C *see* Bongers EMHF *et al*
Boisson C *see* Olschwang S *et al*
Bom SJH *see* Fransen E *et al*
Bonaïti-Pellié C *see* Chompret A *et al*
Bonaglia M *see* Rossi E *et al*
Bonaventure J *see* Francannet C *et al*
Bond J *see* Hampshire DJ *et al*
Bondurand N *see* Pingault V *et al*
Bongers EMHF *et al.* The small patella syndrome: description of five cases from three families and examination of possible allelism with familial patella aplasia-hypoplasia and nail-patella syndrome: *letter*, **209**
Bonneau D *see* Giraudeau F *et al*
Bonnefont J-P *see* Amiel J *et al*
see Imessaoudene B *et al*
Bonne-Tamir B *see* Ben-Yosef T *et al*
Book reviews Lander E, Page D, Lifton R, *editors*. Annual review of genomics and human genetics, **279**
Sankoff D, Nadeau JH, *editors*. Comparative genomics, **807**
Tanzi RE, Parsons AB. Decoding darkness - the search for the genetic causes of Alzheimer's disease, **894**
Verlinsky Y, Kuliev A, *editors*. An atlas of preimplantation genetic diagnosis, **423**
Borck G *et al.* Molecular cytogenetic characterisation of a complex 46,XY,t(7;8;11;13) chromosome rearrangement in a patient with Moebius syndrome: *letter*, **117**
Borg A *see* Bergthorsson JT *et al*
Borgatti R *see* Rossi E *et al*
Borgione E *et al.* How microsatellite analysis can be exploited for subtelomeric chromosomal rearrangement analysis in mental retardation: *letter*, **e1**
Børglum AD *see* Andersen PS *et al*
see Mogensen J *et al*
Borrego S *see* Li Y *et al*
Bost M *see* Faivre L *et al*
Boucher CA *see* Ross NLJ *et al*
Boucher CA *et al.* Breakpoint analysis of Turner patients with partial Xp deletions: implications for the lymphoedema gene location, **591**
Bougeard G *et al.* Detection of 11 germline inactivating *TP53* mutations and absence of *TP63* and *HCHK2* mutations in 17 French families with Li-Fraumeni or Li-Fraumeni-like syndrome: *letter*, **253**
Bourdon V *et al.* Evidence of somatic mosaicism for a *MECP2* mutation in females with Rett syndrome: diagnostic implications: *letter*, **867**
Bourgeron T *see* Quintana-Murci L *et al*
Boutouyrie P *et al.* Non-invasive evaluation of arterial involvement in patients affected with Fabry disease: *letter*, **629**
Bramsen I *see* Henneman L *et al*
Brancolini V *see* Vitale E *et al*
Brayne C *see* Taylor AE *et al*
Breen C *see* McEntagart M *et al*
Brennan P *et al.* Congenital diaphragmatic hernia and interstitial deletion of chromosome 3: *letter*, **556**
Breuning M *see* Maat-Kievit A *et al*
Breuning MH *see* Wagner A *et al*
Briard M-L *see* Edery P *et al*
Brink PA *see* Andersen PS *et al*
Bröcker-Vriends AHJT *see* Wagner A *et al*
Brody LC *see* Eng C *et al*
Brondum-Nielsen K *see* Borck G *et al*
Brook JD *see* Akrami SM *et al*
Brown R *see* Ho LW *et al*
Brugières L *see* Chompret A *et al*
Bruk I *see* Kwok JBJ *et al*
Brundler MA *see* Delozier-Blanchet CD *et al*
Brunoni D *see* Nadal M *et al*
Bryce G *see* Tobias S *et al*
Budarf ML *see* Gong W *et al*
Bugge M *see* Borck G *et al*
Buller RE *see* Mahavni V *et al*
Bulman DE *see* Scoggan KA *et al*
Bundgaard H *see* Andersen PS *et al*
Burden AD *see* Veal CD *et al*
Burger J *see* Vanita *et al*
Butler J *see* Whittington JE *et al*
Buzhievskaya TI *see* Röthlisberger B *et al*
Byers PH *see* Pace JM *et al*
Cabanas R *see* Amaral MD *et al*
Cabrol D *see* Viot G *et al*
Caethoven G *see* van Laer L *et al*
Cairns N *see* Taylor AE *et al*
Cali A *see* Cerruti Mainardi P *et al*
Calvas P *see* Tuffery-Giraud S *et al*
Carey A *see* McEntagart M *et al*
Carles S *see* Tuffery-Giraud S *et al*
Carracedo A *see* Ruiz-Ponte C *et al*
Carrozzo R *see* Rossi E *et al*
Carter N *see* Boardman JP *et al*
Carter ND *see* Syrris P *et al*
Casals T *see* Amaral MD *et al*
Caselli D *see* Rossi E *et al*
Casey B *see* Vitale E *et al*
Castellan C *see* Rossi E *et al*
Castellani C *et al.* Analysis of the entire coding region of the cystic fibrosis transmembrane regulator gene in neonatal hypertrypsinaemia with normal sweat test: *letter*, **202**
Castiglia L *see* Borgione E *et al*
see Rossi E *et al*
Castro M *see* Giannotti A *et al*
Caulder MS *see* Erickson RP *et al*
Caux-Moncoutier V *see* Gad S *et al*
Cavalier L *see* Tuffery-Giraud S *et al*
Cavani S *see* Cerruti Mainardi P *et al*
Cazeneuve C *see* Amaral MD *et al*
Ceccherini I *see* Pingault V *et al*
Cerosaletti KM *see* Hiel JAP *et al*
Cerruti Mainardi P *et al.* Clinical and molecular characterisation of 80 patients with 5p deletion: genotype-phenotype correlation, **151**
Chabrol B *see* de Lonlay P *et al*
Chadwick RB *et al.* Hereditary and somatic DNA mismatch repair gene mutations in sporadic endometrial carcinoma: *letter*, **461**
Chae CU *see* Vitale E *et al*
Chamberlin GP *see* van Laer L *et al*
Chandra T *see* Scoggan KA *et al*
Chantrel K *see* von Kleist-Retzow J-C *et al*
Charbonnier F *see* Bougeard G *et al*
see Saugier-Veber P *et al*
Chartier-Harlin M-C *see* Lambert J-C *et al*
Chatelain P *see* Faivre L *et al*
Chaze AM *see* Lefort G *et al*
Chelly J *see* Bourdon V *et al*
see Villard L *et al*
Chen BLS *see* Sham MH *et al*
Chen J *see* Liu T *et al*
Chen P *see* Lossie AC *et al*
Chen RW *see* Aaltonen L-M *et al*
Chevillard C *see* Villard L *et al*
Chitayat D *see* Dar P *et al*
Chitty LS *see* Kalache KD *et al*
Chomel J-C *see* Amaral MD *et al*
Chompret A *et al.* Sensitivity and predictive value of criteria for *p53* germline mutation screening: *letter*, **43**
Chong B *see* Hegde MR *et al*
Chou S-P *see* Hwu W-L *et al*
Chouery E *see* Mustapha M *et al*
Chowdhury A *see* Kadakol A *et al*
Chowdhury JR *see* Kadakol A *et al*
Chowdhury NR *see* Kadakol A *et al*
Chowdhury S *see* Kadakol A *et al*
Chretien D *see* von Kleist-Retzow J-C *et al*
Christian SL *see* Ungaro P *et al*
Christiansen M *see* Andersen PS *et al*
see Mogensen J *et al*
Chrzanowska KH *et al.* Atypical clinical picture of the Nijmegen breakage syndrome associated with developmental abnormalities of the brain: *letter*, **e3**
Chu CE *see* Flintoff KJ *et al*
Cinti R *et al.* Molecular characterisation of a supernumerary ring chromosome in a patient with VATER association: *letter*, **e6**
Claes E *see* Welkenhuysen M *et al*
Clark AT *see* Dee SL *et al*
Clarke D *see* Whittington JE *et al*
Claustres M *see* Amaral MD *et al*
see Tuffery-Giraud S *et al*
Clavel C *see* Amaral MD *et al*
Clayton-Smith J *see* Watson P *et al*
Clementi R *et al.* Six novel mutations in the *PRF1* gene in children with haemophagocytic lymphohistiocytosis: *letter*, **643**
Clough RL *see* Veal CD *et al*
Coates J *see* Lambert J-C *et al*
Cohen-Tanugi A *see* Francannet C *et al*
Cole TR *see* Baker L *et al*
Colgan J *see* Tobias S *et al*
Colistro F *see* Giannotti A *et al*
Collins J *see* Gong W *et al*
Concannon P *see* Hiel JAP *et al*
Conde R *see* Ruiz-Ponte C *et al*

Conter V *see* Clementi R *et al*
Cooke A *see* Dundar M *et al*
see Tobias S *et al*
Cordier MP *see* Faivre L *et al*
Corfield VA *see* Andersen PS *et al*
Cormier-Daire V *see* de Lonlay P *et al*
see Faivre L *et al*
see Huber C *et al*
see Huber C *et al*
see Imessaoudene B *et al*
see von Kleist-Retzow J-C *et al*
Cornelisse CJ *see* Wagner A *et al*
Costello JF, Plass C. Methylation matters: *review*, **285**
Coubes P *see* Tuffery-Giraud S *et al*
Coucke P *see* van Laer L *et al*
Coucourde G *see* Cerruti Mainardi P *et al*
Coulter-Mackie MB *et al*. Multiple metachromatic leucodystrophy alleles in an unaffected subject: a case of dispermic chimaerism, **e15**
Coupry I *et al*. Leucodystrophy and oculocutaneous albinism in a child with an 11q14 deletion, **35**
Cousin J *see* Faivre L *et al*
Cox LL *see* Kayserili H *et al*
Cox TC *see* Kayserili H *et al*
Cox TM *see* Kelly AL *et al*
Cremers CWRJ *see* Fransen E *et al*
Cremers FPM *see* Fransen E *et al*
Croaker GD *see* Brennan P *et al*
Croquette MF *see* Giraudeau F *et al*
Crow TJ *see* Ross NLJ *et al*
Crow Y *see* Hampshire DJ *et al*
Crow YJ *see* Henwood J *et al*
Cucci RA *see* van Laer L *et al*
Cummings A *see* Lambert J-C *et al*
Cummings SA *see* Fackenthal JD *et al*
Cuppens H *see* Amaral MD *et al*
Cusin V *see* Huber C *et al*

Dadoune JP *see* Siffroi JP *et al*
Dagenais SL *see* Erickson RP *et al*
Dagna Bricarelli F *see* Cerruti Mainardi P *et al*
see Rossi E *et al*
Dagoneau N *see* Huber C *et al*
Dahl H-HM *see* Houseman MJ *et al*
Dallapiccola B *see* Giannotti A *et al*
D'Amelio P *see* Pescarmona GP *et al*
Danesino C *see* Clementi R *et al*
see Maraschio P *et al*
see Rossi E *et al*
Dapena J *see* Amaral MD *et al*
Dar P *et al*. Potential mapping of corneal dermoids to Xq24-qter: *letter*, **719**
David A *see* Giraudeau F *et al*
Dearlove AM *see* Reid E *et al*
de Blois M-C *see* de Leersnyder H *et al*
de Boeck K *see* Amaral MD *et al*
Debré M *see* Edery P *et al*.
Declau F *see* Fransen E *et al*
Decruyenaere M *see* Welkenhuysen M *et al*
Dee SL *et al*. A case of ring chromosome 2 with growth retardation, mild dysmorphism, and microdeletion of 2p detected using FISH, **e32**
de Fusco C *see* Clementi R *et al*
de Kok YJM *see* Fransen E *et al*
del Castillo I *see* Moreno-Pelayo MA *et al*
de la Chapelle A *see* Chadwick RB *et al*
Delague V *see* Mustapha M *et al*
Delattre O *see* Bougeard G *et al*
de Leersnyder H *et al*. β_1-adrenergic antagonists improve sleep and behavioural disturbances in a circadian disorder, Smith-Magenis syndrome, **586**
de Leeuw WJF *see* Wagner A *et al*
Delobel B *see* Giraudeau F *et al*
de Lonlay P *et al*. A broad spectrum of clinical presentations in congenital disorders of glycosylation I: a series of 26 cases, **14**
Delozier-Blanchet CD *et al*. Diaphragmatic hernia, hydrocephalus, and cardiac malformations in four pregnancies of a non-consanguineous couple: *letter*, **269**
Demaille J *see* Lefort G *et al*
Denayer L *see* Welkenhuysen M *et al*
Denton M *see* Dollfus H *et al*
Depetris D *see* Mignon-Ravix C *et al*
de Rienzo A *see* Vitale E *et al*
der Kaloustian VM *see* Lemyre E *et al*
des Georges M *see* Amaral MD *et al*
Devilee P *see* Eng C *et al*
Devoto M *see* Vitale E *et al*
Devriendt K *see* Witters I *et al*
de Vries BBA *et al*. Clinical studies on submicroscopic subtelomeric rearrangements: a checklist, **145**
Submicroscopic subtelomeric 1qter deletions: a recognisable phenotype?: *letter*, **175**
Deymeer F *see* Akbas F *et al*
Di WL *see* Houseman MJ *et al*
Diaw L *see* Siwarski D *et al*
Dietz H *see* Gong W *et al*
Dietze I *see* Fritz B *et al*
Dietzmann K *see* Kirches E *et al*
Digilio MC *see* Giannotti A *et al*
Dinulos MB *see* Parisi MA *et al*
di Rocco M *see* Rossi E *et al*
Dobyns WB *see* Ungaro P *et al*
Dollfus H *et al*. Identification of a new *TWIST* mutation (7p21) with variable eyelid manifestations supports locus homogeneity of BPES at 3q22: *letter*, **470**
Dommergues M *see* Amiel J *et al*
Dong HJ *see* Lossie AC *et al*
Dörk T *see* Amaral MD *et al*
see Hughes D *et al*
see Ockenga J *et al*
Doudounakis S *see* Amaral MD *et al*
Downes SM *see* Payne AM *et al*
Downing S *see* O'Shaughnessy KM *et al*
Downs CA *see* Erickson RP *et al*
Driscoll DA *see* Gong W *et al*
Driscoll DJ *see* Lossie AC *et al*
Drouin V *see* de Lonlay P *et al*
Drouot N *see* Saugier-Veber P *et al*
Duisterhof M *et al*. Psychological studies in Huntington's disease: making up the balance: *letter*, **852**
Duncan AMV *see* Lemyre E *et al*
Dundar M *et al*. A novel acropectoral syndrome maps to chromosome 7q36, **304**
Dupont JM *see* Viot G *et al*

Easton D *see* Taylor AE *et al*
Ebeling T *see* Kytölä S *et al*
Ebenezer N *see* Talks SJ *et al*
Eccles D *et al*. A novel 3' mutation in the *APC* gene in a family presenting with a desmoid tumour: *letter*, **861**
Echenne B *see* Tuffery-Giraud S *et al*
Edery P *et al*. B cell immunodeficiency, distal limb abnormalities, and urogenital malformations in a three generation family: a novel autosomal dominant syndrome?: *letter*, **488**
Egeblad H *see* Mogensen J *et al*
Eggermann K *see* Albrecht B *et al*
see Eggermann T *et al*
Eggermann T *see* Albrecht B *et al*
Eggermann T *et al*. Identification of interstitial maternal uniparental disomy (UPD) (14) and complete maternal UPD(20) in a cohort of growth retarded patients, **86**
Ehrlich M *et al*. High frequencies of ICF syndrome-like pericentromeric heterochromatin decondensation and breakage in chromosome 1 in a chorionic villus sample : *letter*, **882**
Eisenbruch M *see* Meiser B *et al*
Ejlertsen B *see* Bergthorsson JT *et al*
Elghezal H *see* Mégarbané A *et al*
Eller P *et al*. Wolfram syndrome: a clinical and molecular genetic analysis: *letter*, **e37**
Ellis LA *see* Houseman MJ *et al*
Emanuel BS *see* Gong W *et al*
Eng C *see* Bender BU *et al*
see Fackenthal JD *et al*
see Parisi MA *et al*
see Reardon W *et al*
Eng C *et al*. Interpreting epidemiological research: blinded comparison of methods used to estimate the prevalence of inherited mutations in *BRCA1* , **824**
Erginel-Unaltuna N *see* Akbas F *et al*
Erickson RP *et al*. Clinical heterogeneity in lymphoedema-distichiasis with *FOXC2* truncating mutations, **761**
Escayg A *see* Reid E *et al*
Esiri MM *see* Taylor AE *et al*
Estivill X *see* Amaral MD *et al*
see Nadal M *et al*
Evans C *see* Fokstuen S *et al*
Evans DGR *see* Baser ME *et al*
see Hopwood P *et al*
see Verma L *et al*
Evans DGR *et al*. Science or black magic?: *letter*, **e16**
Evans JG *see* Taylor AE *et al*
Evers-Kiebooms G *see* Welkenhuysen M *et al*
Extermann P *see* Delozier-Blanchet CD *et al*

Fackenthal JD *et al*. Male breast cancer in Cowden syndrome patients with germline *PTEN* mutations, **159**
Fahsold R *see* Luijten M *et al*
Failla P *see* Borgione E *et al*
Faivre L *see* Viot G *et al*
Faivre L *et al*. Acromicric dysplasia: long term outcome and evidence of autosomal dominant inheritance, **745**
Fallet-Bianco C *see* Viot G *et al*
Fannin TF *see* McConnell RS *et al*

Fantes JA *see* Ungaro P *et al*
FAP Collaborative Research Group *see* Michie *et al*
Faravelli F *see* Rossi E *et al*
Farinha CM *see* Amaral MD *et al*
Farmer G *see* Tobias S *et al*
Fawkner M *see* Hegde MR *et al*
Feingold J *see* Amiel J *et al*
see Gabolde M *et al*
Fenger K *see* Bergthorsson JT *et al*
Férec C *see* Mura C *et al*
see Amaral MD *et al*
Ferreira JC *see* Dar P *et al*
Ferreira P *see* Coulter-Mackie MB *et al*
Ferrero GB *see* Vitale E *et al*
Ferretti F *see* Giannotti A *et al*
Ferry B *see* Veal CD *et al*
Feunteun J *see* Chompret A *et al*
Fichera M *see* Borgione E *et al*
Filocamo M *et al*. A simple non-isotopic method to show pitfalls during mutation analysis of the glucocerebrosidase gene: *letter*, **e34**
Finidori G *see* Faivre L *et al*
Fink JK *see* Rainier S *et al*
Finkenstedt G *see* Eller P *et al*
Finnilä S *et al*. Increased risk of sensorineural hearing loss and migraine in patients with a rare mitochondrial DNA variant 4336A>G in $tRNA^{Gln}$: *letter*, **400**
Firatli E *see* Zhang Y *et al*
Fischer A *see* Edery P *et al*
Fischer H *see* Haberlandt E *et al*
Fitzpatrick J *see* Grunebaum E *et al*
Flaman J-M *see* Bougeard G *et al*
Flint J *see* de Vries BBA *et al*
Flinter FA *see* King K *et al*
Flintoff KJ *et al*. Submicroscopic deletions of the *APC* gene: a frequent cause of familial adenomatous polyposis that may be overlooked by conventional mutation scanning: *letter*, **129**
Flint J *see* de Vries BBA *et al*
Flothmann K *see* van Laer L *et al*
Flynn DM *see* Kelly AL *et al*
Fodde R *see* Wagner A *et al*
Föger B *see* Eller P *et al*
Fokstuen S *et al*. Eight years' experience of direct molecular testing for myotonic dystrophy in Wales: *letter*, **e42**
Presymptomatic testing in myotonic dystrophy: genetic counselling approaches : *letter*, **846**
Fontès M *see* Villard L *et al*
Foulkes WD *see* Kasprzak L *et al*
Francannet C *et al*. Genotype-phenotype correlation in hereditary multiple exostoses, **430**
Frank TS *see* Eng C *et al*
Fransen E *see* van Laer L *et al*
Fransen E *et al*. A common ancestor for *COCH* related cochleovestibular (DFNA9) patients in Belgium and The Netherlands bearing the P51S mutation: *letter*, **61**
Frébourg T *see* Bougeard G *et al*
see Saugier-Veber P *et al*
Fricker J-P *see* Bougeard G *et al*
Friedman JM *see* Baser ME *et al*
Friedman TB *see* Ben-Yosef T *et al*
Fritz B *et al*. A supernumerary marker chromosome with a neocentromere derived from 5p14→pter: *letter*, **559**
Fryns J-P *see* Syrrou M and Fryns J-P
see Witters I *et al*
Fu B *see* O'Shaughnessy KM *et al*
Fu M *see* Sham MH *et al*
Fujimoto S *see* Muroya K *et al*
Fukushima Y *see* Muroya K *et al*
Fusch C *see* Eggermann T *et al*

Gabolde M *et al*. The mannose binding lectin gene influences the severity of chronic liver disease in cystic fibrosis, **310**
Gaboury L *see* Kasprzak L *et al*
Gabreëls FJM *see* Hiel JAP *et al*
Gabriel BM *see* de Lonlay P *et al*
Gad S *et al*. Identification of a large rearrangement of the *BRCA1* gene using colour bar code on combed DNA in an American breast/ovarian cancer family previously studied by direct sequencing: *letter*, **388**
Gaff C *et al*. Oncology nurse training in cancer genetics: *letter*, **691**
Galesi O *see* Borgione E *et al*
see Rossi E *et al*
Gambarara M *see* Giannotti A *et al*
Gander R *see* Eller P *et al*
Ganev V *see* Horvath A *et al*
Gartner S *see* Amaral MD *et al*
Gassner I *see* Janecke AR *et al*
Gatti R *see* Filocamo M *et al*
Gauthier G *see* Monnier N *et al*
Gerbitz K-D *see* Jaksch M *et al*
Germain DP *see* Boutouyrie P *et al*
Germain DP. Pseudoxanthoma elasticum: evidence for the existence of a pseudogene highly homologous to the *ABCC6* gene: *letter*, 457
Ghassibeh M *see* Mustapha M *et al*
Ghosh SS *see* Kadakol A *et al*
Giangaspero F *see* Steigerwald K *et al*
Giannotti A *et al*. Coeliac disease in Williams syndrome, **767**
Gille JJP *see* Henneman L *et al*
Gimelli G *see* Cinti R *et al*
Girardet A *see* Lefort G *et al*
Giraudeau F *et al*. Use of a set of highly polymorphic minisatellite probes for the identification of cryptic 1p36.3 deletions in a large collection of patients with idiopathic mental retardation: *letter*, **121**
Girodon-Boulandet E *see* Amaral MD *et al*
Gismondi V *see* Montera M *et al*
see Steigerwald K *et al*
Giudice ML *see* Borgione E *et al*
Glaser T *see* Vincent A *et al*
Glover TW *see* Erickson RP *et al*
Goizet C *see* Coupry I *et al*
Goldmuntz E *see* Gong W *et al*
Goldstein D *see* Meiser B *et al*
Gollnick HPM *see* Zouboulis CC *et al*
Gong W *et al*. Mutation analysis of *TBX1* in non-deleted patients with features of DGS/VCFS or isolated cardiovascular defects, **e45**
Goodman FR *see* Dundar M *et al*
Goodship JA *see* Tompson SWJ *et al*
Goossens M *see* Amaral MD *et al*
see Pingault V *et al*
Gordon TM *see* Dundar M *et al*
Gore M *see* Hallowell N *et al*
Gorry MC *see* Zhang Y *et al*
Gosset P *see* Mégarbané A *et al*
Gottlieb S *see* Gong W *et al*
Goudev A *see* Horvath A *et al*
Gout J-P *see* Monnier N *et al*
Graham C *see* Hughes D *et al*
Graham JS *see* Steigerwald K *et al*
Greco D *see* Rossi E *et al*
Green AJ *see* Willatt L *et al*
Green PM *see* King K *et al*
Greenamyre JT *see* Simon DK *et al*
Green AJ *see* McEntagart M *et al*
Gregersen N *see* Mogensen J *et al*
Gregorek H *see* Chrzanowska KH *et al*
Gregory-Evans CY *see* Talks SJ *et al*
see Tarttelin EE *et al*
Gregory-Evans K *see* Talks SJ *et al*
see Tarttelin EE *et al*
Griffith AJ. Genetic analysis of the connexin-26 M34T variant: *letter*, **e24**
Griscelli C *see* Edery P *et al*
Groden J *see* Steigerwald K *et al*
Grönberg H *see* Liu T *et al*
Gross SJ *see* Dar P *et al*
Grossmann JG *see* Evans DGR *et al*
Grumpelt U *see* Tönnies H *et al*
Grunebaum E *et al*. A missense mutation in the *SEDL* gene results in delayed onset of X linked spondyloepiphyseal dysplasia in a large pedigree: *letter*, **409**
Grunfeld J-P *see* Boutouyrie P *et al*
Guala A *see* Maraschio P *et al*
Guanti G *see* Montera M *et al*
Guilloud-Bataille M *see* Gabolde M *et al*
Guittard C *see* Tuffery-Giraud S *et al*

Haaf T *see* Borck G *et al*
Haberlandt E *et al*. Split hand/split foot malformation associated with sensorineural deafness, inner and middle ear malformation, hypodontia, congenital vertical talus, and deletion of eight microsatellite markers in 7q21.1-q21.3: *letter*, **405**
Haddad BR *see* Stratakis CA *et al*
Hagerman PJ *see* Tassone F *et al*
Hagerman RJ *see* Tassone F *et al*
Hahn AF *see* Scoggan KA *et al*
Hall CM *see* Offiah AC *et al*
Hallowell N *et al*. Surveillance or surgery? A description of the factors that influence high risk premenopausal women's decisions about prophylactic oophorectomy: *letter*, **683**
Hallowell U *see* Hallowell N *et al*
Haltia T *see* Nopola-Hemmi J *et al*
Hamel BCJ *see* Bongers EMHF *et al*
Hampel H *see* Chadwick RB *et al*
Hampshire DJ *et al*. Kufor-Rakeb syndrome, pallido-pyramidal degeneration with supranuclear upgaze paresis and dementia, maps to 1p36, **680**
Hannula K *et al*. Do patients with maternal uniparental disomy for chromosome 7 have a distinct mild Silver-Russell phenotype?: *letter*, **273**
Harboe TL *see* Bergthorsson JT *et al*

Hardt T *see* Borck G *et al*
Harper J *see* Barker K *et al*
Harper P. Cyril Clarke, *Journal of Medical Genetics*, and the foundation of clinical genetics, **283**
Harper PS *see* Fokstuen S *et al*
see Fokstuen S *et al*
see Simpson SA *et al*
Harris JM *see* Lambert J-C *et al*
Hart PS *see* Zhang Y *et al*
Hart TC *see* Zhang Y *et al*
Hartikainen P *see* Finnilä S *et al*
Harvey J *see* Eccles D *et al*
Hasegawa T *see* Muroya K *et al*
Hasegawa Y *see* Muroya K *et al*
Hasnain SS *see* Evans DGR *et al*
Hata A *see* Yabe I *et al*
Hauck Y *see* Giraudeau F *et al*
Haukkala A *see* Aktan-Collan K *et al*
Havndrup O *see* Andersen PS *et al*
Heath M *see* Brennan P *et al*
Heathcote G *see* Vincent A *et al*
Hedera P *see* Rainier S *et al*
Hegde MR *et al.* Microdeletion in the *FMR-1* gene: an apparent null allele using routine clinical PCR amplification: *letter*, **624**
Heiman-Patterson T *see* Rainier S *et al*
Heinz-Erian P *see* Haberlandt E *et al*
Hendriks Y *see* Wagner A *et al*
Heng CK *see* Liu Y *et al*
Henneman L *et al.* Participation in preconceptional carrier couple screening: characteristics, attitudes, and knowledge of both partners: *letter*, **695**
Hennies H-C *see* Tönnies H *et al*
Henske EP *see* Khare L *et al*
Henwood J *et al.* A region of homozygosity within 22q11.2 associated with congenital heart disease: recessive DiGeorge/velocardiofacial syndrome?: *letter*, **533**
Héon E *see* Vincent A *et al*
Herman G *see* Erickson RP *et al*
Hermsen KJM *see* Zusterzeel PLM *et al*
Hernández-Calvin FJ *see* Moreno-Pelayo MA *et al*
Herraiz C *see* Moreno-Pelayo MA *et al*
Herrera D *see* Ehrlich M *et al*
Hiel JAP *et al.* Nijmegen breakage syndrome in a Dutch patient not resulting from a defect in *NBS1*: *letter*, **e19**
Hill S *see* Stratakis CA *et al*
Hirst-Stadlmann A *see* Haberlandt E *et al*
Hitchins M *see* Monk D *et al*
Hitchins MP *et al.* Silver-Russell syndrome: a dissection of the genetic aetiology and candidate chromosomal regions: *review*, **810**
Ho LW *et al.* Wild type huntingtin reduces the cellular toxicity of mutant huntingtin in mammalian cell models of Huntington's disease, **450**
Hoffman-Radvanyi H *see* Amiel J *et al*
Höög A *see* Kytölä S *et al*
Hofstra R *see* Wagner A *et al*
Hoganson GE *see* Tahmaz FE *et al*
Hokezu Y *see* Yabe I *et al*
Holder SE *see* Boardman JP *et al*
Holland AJ *see* Whittington JE *et al*
Holloway S *see* Dundar M *et al*
Holmberg E *see* Liu T *et al*
Holmberg C *see* Hannula K *et al*
Holmes A *see* MacDermot KD *et al*
see MacDermot KD *et al*
Homfray T *see* de Vries BBA *et al*
Hong S *see* Liu Y *et al*
Hopwood P *et al.* Risk perception and cancer worry: an exploratory study of the impact of genetic risk counselling in women with a family history of breast cancer: *letter*, **139**
Horn D *et al.* Anauxetic dysplasia, a spondylometaepiphyseal dysplasia with extreme dwarfism: *letter*, **262**
Horvath A *et al.* High frequency of the *ApoB*-100 R3500Q mutation in Bulgarian hypercholesterolaemic subjects: *letter*, **536**
Horvath R *see* Jaksch M *et al*
Houlston R *see* Barker K *et al*
Houlston RS *see* Tomlinson IPM *et al*
Houseman M *see* van Laer L *et al*
Houseman MJ *see* Kelsell DP *et al*
Houseman MJ *et al.* Genetic analysis of the connexin-26 M34T variant: identification of genotype M34T/M34T segregating with mild-moderate non-syndromic sensorineural hearing loss, **20**
Howell A *see* Evans DGR *et al*
Howell RT *see* Baker L *et al*
Howell A *see* Hopwood P *et al*
Hrafnkelsson J *et al.* Familial non-medullary thyroid cancer in Iceland: *letter*, **189**
Hu Y-H *see* Hwu W-L *et al*
Huber C *et al. SHOX* point mutations in dyschondrosteosis: *letter*, **323**
Sulphate transporter gene mutations in apparently isolated club foot: *letter*, **191**
Hubert D *see* Amaral MD *et al*
see Gabolde M *et al*
Hudgins L *see* Parisi MA *et al*
Huggare J *see* Wong FK *et al*
Hughes AE *see* McConnell RS *et al*
Hughes D *et al.* Mutation and haplotype analysis of the *CFTR* gene in atypically mild cystic fibrosis patients from Northern Ireland: *letter*, **136**
Hughes HE *see* Baker L *et al*
Hugot J-P *see* Pingault V *et al*
Hulsebos TJM *see* Luijten M *et al*
Hunt DM *see* Payne AM *et al*
Huppi K *see* Siwarski D *et al*
Hurst JA *see* de Vries BBA *et al*
Hutchin TP *et al.* Prevalence of mitochondrial DNA mutations in childhood/congenital onset non-syndromal sensorineural hearing impairment, **229**
Hutson A *see* Lossie AC *et al*
Huygen PLM *see* Fransen E *et al*
Hwu W-L *et al.* A Chinese adult onset type II citrullinaemia patient with 851del4/1638ins23 mutations in the *SLC25A13* gene: *letter*, **e23**
Hykin P *see* Talks SJ *et al*

Ibañez A *see* Lorda-Sanchez I *et al*
Ibi T *see* Sahashi K *et al*
Iguchi A *see* Kobayashi I *et al*
Imamura A *et al.* Temperature sensitive acyl-CoA oxidase import in group A peroxisome biogenesis disorders: *letter*, **871**
Imessaoudene B *et al. MECP2* mutation in non-fatal, non-progressive encephalopathy in a male, **171**
Inoue I *see* Yabe I *et al*
Isaia GC *see* Pescarmona GP *et al*
Ishikawa N *see* Kobayashi I *et al*
Isotani H *see* Muroya K *et al*
Ito Y *see* Muroya K *et al*
Iwata Y *see* Muroya K *et al*
Iwatsubo T *see* Lambert J-C *et al*

Jacobs I *see* Hallowell N *et al*
Jaeken J *see* de Lonlay P *et al*
Jaksch M *et al.* Frequency of mitochondrial transfer RNA mutations and deletions in 225 patients presenting with respiratory chain deficiencies, **665**
James V. The importance of good images in using hair to screen for breast cancer: *letter*, **e16**
Janecke AR *see* Haberlandt E *et al*
Janecke AR *et al.* Adducted thumb-club foot syndrome in sibs of a consanguineous Austrian family: *letter*, **265**
Janka G *see* Clementi R *et al*
Jansen RPM *see* van der Luijt RB *et al*
Jaruzelska J *et al.* Mosaicism for 45,X cell line may accentuate the severity of spermatogenic defects in men with AZFc deletion, **798**
Javed AA *see* Dar P *et al*
Jedrzejczak P *see* Jaruzelska J *et al*
Jeffery S *see* Syrris P *et al*
Jóźwiak S *see* Chrzanowska KH *et al*
Johnson CK *see* Chadwick RB *et al*
Johnson S *see* Payne AM *et al*
Johns DR *see* Simon DK *et al*
Johnston H *see* Jones KJ *et al*
Joly G *see* Mégarbané A *et al*
Jónasson JG *see* Hrafnkelsson J *et al*
Jones AB *see* Veal CD *et al*
Jones KJ *et al.* The expanding phenotype of laminin α2 chain (merosin) abnormalities: case series and review: *review*, **649**
Jones MC *see* Erickson RP *et al*
Jonveaux P *see* Bourdon V *et al*
see Bougeard G *et al*
Jouannet P *see* Viot G *et al*
Jouannic J-M *see* Amiel J *et al*
Jouk P-S *see* Pingault V *et al*
Journel H *see* de Lonlay P *et al*
see Giraudeau F *et al*
Judmaier W *see* Haberlandt E *et al*
Jung M *see* Vanita *et al*

Kadakol A *et al.* Interaction of coding region mutations and the Gilbert-type promoter abnormality of the *UGT1A1* gene causes moderate degrees of unconjugated hyperbilirubinaemia and may lead to neonatal kernicterus: *letter*, **244**
Kääriäinen H *see* Aktan-Collan K *et al*
Kagawa S *see* Kanayama H-O *et al*
Kaji M *see* Yorifuji T *et al*
Kalache KD *et al.* Laryngeal atresia, encephalocele, and limb deformities (LEL): a possible new syndrome: *letter*, **420**
Kanafani S *see* Siffroi JP *et al*
Kanavakis E *see* Amaral MD *et al*
Kanayama H-O *et al.* Association of a novel constitutional translocation t(1q;3q) with familial renal cell carcinoma, **165**
Karshelova E *see* Horvath A *et al*

Karsten A *see* Wong FK *et al*
Kasprzak L *et al*. Familial medullary thyroid carcinoma and prominent corneal nerves associated with the germline V804M and V778I mutations on the same allele of *RET*: *letter*, **784**
Kattner E *see* Fritz B *et al*
Kawai M *see* Yorifuji T *et al*
Kawamura N *see* Kobayashi I *et al*
Kayserili H *et al*. Molecular characterisation of a new case of microphthalmia with linear skin defects (MLS): *letter*, **411**
Kedra D *see* Kanayama H-O *et al*
Keitzer R *see* Tönnies H *et al*
Kelly AL *et al*. Classification and genetic features of neonatal haemochromatosis: a study of 27 affected pedigrees and molecular analysis of genes implicated in iron metabolism, **599**
Kelly B *see* McConnell RS *et al*
Kelly DA *see* Kelly AL *et al*
Kelsell DP *see* Houseman MJ *et al*
see Rickard S *et al*
Kelsell DP *et al*. Reply: *letter*, **e24**
Kemperman MH *see* Fransen E *et al*
Kenneson A *see* Hegde MR *et al*
Kentala A *see* Pirinen S *et al*
Kere J *see* Hannula K *et al*
see Nopola-Hemmi J *et al*
see Wong FK *et al*
Kerr B *see* Watson P *et al*
Kerstjens-Frederikse WS *see* Erickson RP *et al*
Khare L *et al*. A novel missense mutation in the GTPase activating protein homology region of *TSC2* in two large families with tuberous sclerosis complex: *letter*, **347**
Kida H *see* Yabe I *et al*
Kikuchi S *see* Komeichi K *et al*
Kiliç G *see* Kayserili H *et al*
Killian A *see* Bougeard G *et al*
Kim J *see* Siwarski D *et al*
Kim SC *see* Mahavni V *et al*
Kindermans C *see* de Leersnyder H *et al*
King K *et al*. Hair roots as the ideal source of mRNA for genetic testing: *letter*, **e20**
King MD *see* McEntagart M *et al*
Kirches E *et al*. Heterogeneous tissue distribution of a mitochondrial DNA polymorphism in heteroplasmic subjects without mitochondrial disorders, **312**
Kirchhoff M *et al*. High resolution comparative genomic hybridisation in clinical cytogenetics, **740**
Kirov S *see* Horvath A *et al*
Kirste G *see* Bender BU *et al*
Kiviluoto T *see* Avizienyte E *et al*
Kjeldsen K *see* Andersen PS *et al*
Klausen S *see* Bergthorsson JT *et al*
Klein ML *see* Tarttelin EE *et al*
Kleinle S *see* Jaksch M *et al*
Kleopa KA *see* Rainier S *et al*
Klersy C *see* Clementi R *et al*
Klopstock T *see* Jaksch M *et al*
Knight SJL *see* de Vries BBA *et al*
see de Vries BBA *et al*
Knoers NVAM *see* Bongers EMHF *et al*
Kobayashi I *et al*. Novel mutations of *FOXP3* in two Japanese patients with immune dysregulation, polyendocrinopathy, enteropathy, X linked syndrome (IPEX): *letter*, **874**
Kobayashi K *see* Hwu W-L *et al*
see Kobayashi I *et al*
Koenig B *see* Bourdon V *et al*
Koillinen H *see* Wong FK *et al*
Komeichi K *et al*. Twenty CAG repeats are sufficient to cause the SCA6 phenotype: *letter*, **e38**
Kondo N *see* Imamura A *et al*
Kooyman MAP *see* Bongers EMHF *et al*
Korcz A *see* Jaruzelska J *et al*
Kordich JJ *see* Steigerwald K *et al*
Korner C *see* de Lonlay P *et al*
Kotecki M *see* Jaruzelska J *et al*
Kotzot D *see* Röthlisberger B *et al*
Kotzot D. Complex and segmental uniparental disomy (UPD): review and lessons from rare chromosomal complements: *review*, **497**
Kotzot D *et al*. Maternal uniparental isodisomy 11q13→qter in a dysmorphic and mentally retarded female with partial trisomy mosaicism 11q13→qter : *letter*, **876**
Kpebe A *see* Villard L *et al*
Krajewska-Walasek M *see* Chrzanowska KH *et al*
Krause G *see* Kirches E *et al*
Kreczy A *see* Janecke AR *et al*
Kretz M *see* de Lonlay P *et al*
Kroos MA *see* Ausems MGEM *et al*
Kudoh J *see* Ben-Yosef T *et al*
Kuismanen S *see* Liu T *et al*
Kumaramanickavel G *see* Dollfus H *et al*
Kunze J *see* Horn D *et al*
Kuroki Y *see* Kanayama H-O *et al*
Kuslich CD *see* Pace JM *et al*
Kwok JBJ *et al*. Mutations in the glycine receptor α1 subunit (*GLRA1*) gene in hereditary hyperekplexia pedigrees: evidence for non-penetrance of mutation Y279C: *letter*, **e17**
Kytölä S *et al*. Founder effect in multiple endocrine neoplasia type 1 (MEN 1) in Finland: *letter*, **185**

Labelle V *see* Villard L *et al*
Lacombe D *see* Bahuau M *et al*
see Coupry I *et al*
see Giraudeau F *et al*
Lafferty A *see* Stratakis CA *et al*
LaFranchi S *see* Beals RK and LaFranchi S
Laganà C *see* Priolo M and Laganà C
Lakhoo K *see* Boardman JP *et al*
Lalloo F *see* Hopwood P *et al*
Laloux B *see* Boutouyrie P *et al*
Lam C-W *et al*. Homoplasmic 3316G→A in the *ND1* gene of the mitochondrial genome: a pathogenic mutation or a neutral polymorphism?: *letter*, **e10**
Lambert J-C *et al*. The −48 CT polymorphism in the presenilin 1 promoter is associated with an increased risk of developing Alzheimer's disease and an increased Aβ load in brain, **353**
Lambiris N *see* Hegde MR *et al*
Lamireau T *see* Bahuau M *et al*
Larsen LA *see* Andersen PS *et al*
Larson O *see* Wong FK *et al*
Larsson C *see* Kanayama H-O *et al*
see Kytölä S *et al*
see Wong FK *et al*
Latif F *see* Webb T and Latif F
Laubenberger J *see* Bender BU *et al*
Launonen V *see* Avizienyte E *et al*
Laurent J *see* de Lonlay P *et al*
Laurent S *see* Boutouyrie P *et al*
Lavinha J *see* Amaral MD *et al*
Le Bail B *see* Bahuau M *et al*
Le Bourhis C *see* Siffroi JP *et al*
Lechleitner M *see* Eller P *et al*
Ledbetter DH *see* Ungaro P *et al*
Le Deist F *see* Edery P etal
Lee DD *see* Reid E *et al*
Lee Y-A *see* Bayoumi R *et al*
Leek JP *see* Henwood J *et al*
Lees M *see* Offiah AC *et al*
Lefebvre S *see* Saugier-Veber P *et al*
Lefort G *et al*. Cytogenetic and molecular study of a jumping translocation in a baby with Dandy-Walker malformation: *letter*, **67**
Le Gac G *see* Mura C *et al*
Legeai-Mallet L *see* Francannet C *et al*
Legius E *see* Witters I *et al*
Lehtovirta M *see* Finnilä S *et al*
Leisti J *see* Kytölä S *et al*
Le Lorc'h M *see* Mégarbané A *et al*
Lemaire F *see* Fransen E *et al*
Le Merrer M *see* de Lonlay P *et al*
see Faivre L *et al*
see Francannet C *et al*
see Huber C *et al*
see Huber C *et al*
Lemmon H *see* Lambert J-C *et al*
Lemort N *see* Pingault V *et al*
Lemyre E *et al*. Stable non-Robertsonian dicentric chromosomes: four new cases and a review: *letter*, **76**
Lenaerts C *see* Gabolde M *et al*
Lench NJ *see* Henwood J *et al*
Lendon C *see* Lambert J-C *et al*
Leppig KA *see* Parisi MA *et al*
Lerone M *see* Cinti R *et al*
Leroy A *see* de Lonlay P *et al*
Lespinasse J *see* Delozier-Blanchet CD *et al*
Levin A *see* Vincent A *et al*
Lévy N *see* Villard L *et al*
Li Y *et al*. Evaluation of the *ELOVL4* gene in families with retinitis pigmentosa linked to the *RP25* locus: *letter*, **478**
Lidove O *see* Boutouyrie P *et al*
Lidral AC *see* Erickson RP *et al*
Liechti-Gallati S *see* Jaksch M *et al*
Ligtenberg M *see* Hiel JAP *et al*
Limacher J-M *see* Bougeard G *et al*
Lindblom U *see* Liu T *et al*
Linder-Aronson S *see* Wong FK *et al*
Lindhout D *see* Wagner A *et al*
Linnemann K *see* Eggermann T *et al*
Lipsanen-Nyman M *see* Hannula K *et al*
Liu HX *et al*. Interstitial deletion of 3p22.2-p24.2: the first reported case: *letter*, **349**
Liu T *et al*. Screening families with endometrial and colorectal cancers for germline mutations: *letter*, **e29**
Liu Y *et al*. Fibrinogen genotypes (α and β) are associated with plasma fibrinogen levels in Chinese, **e30**

Lochmuller H *see* Jaksch M *et al*
Löffler J *see* Haberlandt E *et al*
Loiseau HA *see* Giraudeau F *et al*
Loiselet J *see* Mustapha M *et al*
Longy M *see* Bougeard G *et al*
Lopes AI *see* Seixas S *et al*
Lopes B *see* Amaral MD *et al*
Lorda-Sanchez I *et al*. Does Bardet-Biedl syndrome have a characteristic face?: *letter*, **e14**
Losekoot M *see* Maat-Kievit A *et al*
Lossie AC *et al*. Distinct phenotypes distinguish the molecular classes of Angelman syndrome, **834**
Louis DN *see* Blakley P *et al*
Lovę DR *see* Hegde MR *et al*
Low PS *see* Liu Y *et al*
Lowenheim M *see* Kadakol A *et al*
Lowy AM *see* Steigerwald K *et al*
Ludman MD *see* Coulter-Mackie MB *et al*
Lui VCH *see* Sham MH *et al*
Lui W-O *see* Kanayama H-O *et al*
Luijten M *et al*. Limited contribution of interchromosomal gene conversion to *NF1* gene mutation: *letter*, **481**
Lunardi J *see* Monnier N *et al*
Lundgren T *see* Zhang Y *et al*
Lundsteen C *see* Kirchhoff M *et al*
Lunt PW *see* Kelly AL *et al*
Lyon G *see* Imessaoudene B *et al*
Lyonnet S *see* Amiel J and Lyonnet S
see Bahuau M *et al*
see Edery P *et al*
see Imessaoudene B *et al*
see Viot G *et al*

Maat-Kievit AA *see* van den Boer-van den Berg HM and Maat-Kievit AA
Maat-Kievit A *et al*. New problems in testing for Huntington's disease: the issue of intermediate and reduced penetrance alleles: *letter*, **e12**
MacArdle B *see* Rickard S *et al*
MacCollin M *see* Blakley P *et al*
McConnell RS *et al*. Autosomal dominant polycystic kidney disease unlinked to the *PKD1* and *PKD2* loci presenting as familial cerebral aneurysm: *letter*, **238**
MacDermot KD *et al*. Anderson-Fabry disease: clinical manifestations and impact of disease in a cohort of 60 obligate carrier females: *letter*, **769**
Anderson-Fabry disease: clinical manifestations and impact of disease in a cohort of 98 hemizygous males, **750**
McDonald M *see* Erickson RP *et al*
McDonald-McGinn DM *see* Gong W *et al*
Macdonald F *see* Verma L *et al*
McEntagart M *et al*. Molecular characterisation of a proximal chromosome 18q deletion: *letter*, **128**
McGaughran J *see* Hegde MR *et al*
McGaughran JM *see* Liu HX *et al*
McGuirt WT *see* Fransen E *et al*
Mackay J *see* Hallowell N *et al*
McKenna WM *see* Syrris P *et al*
MacKenzie JJ *see* Grunebaum E *et al*
McKeown C *see* de Vries BBA *et al*
McKiernan PJ *see* Kelly AL *et al*
McKinstry CS *see* McConnell RS *et al*
McKusick VA. Persisting memories of Cyril Clarke in Baltimore, **284**
McQuaid S *see* McEntagart M *et al*
Mahavni V *et al*. The androgen receptor and DXS15-134 markers show a high rate of discordance for germline X chromosome inactivation: *letter*, **474**
Maher ER. *JMG* Online: exploiting the potential of electronic publication and manuscript submission: *editorial*, **809**
Maher ER *see* Verma L *et al*
Majamaa K *see* Finnilä S *et al*
Mäkitie AA *see* Aaltonen L-M *et al*
Malcolm S *see* de Vries BBA *et al*
see Ungaro P *et al*
Maltby EL *see* Shannon NL *et al*
Mamada M *see* Yorifuji T *et al*
Mann DMA *see* Lambert J-C *et al*
Mannermaa A *see* Finnilä S *et al*
Maraschio P *et al*. A novel mutation and novel features in Nijmegen breakage syndrome: *letter*, **113**
Marchese C *see* Montera M *et al*
Marcos I *see* Li Y *et al*
Mareni C *see* Montera M *et al*
Marks JJ *see* Zhang Y *et al*
Maroteaux P *see* Faivre L *et al*
Marsh DJ *see* Fackenthal JD *et al*
Marteau TM *see* Michie S *et al*
Martin C *see* Bougeard G *et al*
Martin DM *et al*. Ocular malformations, postaxial polydactyly, and delayed intramembranous ossification: a new autosomal dominant condition: *letter*, **547**
Martinez A *see* Barker K *et al*
Mason S *see* Veal CD *et al*
Mastella G *see* Castellani C *et al*
Masturzo B *see* Kalache KD *et al*
Mathieu M *see* Huber C *et al*
Matsell D *see* Rupar CA *et al*
Matsuo N *see* Ogata T *et al*
Mattei MG *see* Giraudeau F *et al*
Mattei M-G *see* Mignon-Ravix C *et al*
Matthijs G *see* de Lonlay P *et al*
Maw M *see* Dollfus H *et al*
Mawrin C *see* Kirches E *et al*
Maxwell M *see* Ho LW *et al*
Mazzotti R *see* Filocamo M *et al*
Meck JM *see* Stratakis CA *et al*
Mecklin J-P *see* Aktan-Collan K *et al*
Meek JE *see* Chadwick RB *et al*
Mégarbané A *see* Mustapha M *et al*
Mégarbané A *et al*. Pure partial 7p trisomy including the *TWIST*, *HOXA*, and *GLI3* genes: *letter*, **178**
Meijers-Heijboer EJ *see* Wagner A *et al*
Meiser B *et al*. Cultural aspects of cancer genetics: setting a research agenda: *review*, **425**
Meisler MH *see* Reid E *et al*
Melaragno MI *see* Nadal M *et al*
Mercier A-Y *see* Mura C *et al*
Meredith L *see* Fokstuen S *et al*
Mergenthaler S *see* Albrecht B *et al*
see Eggermann T *et al*
Michael M *see* Kirches E *et al*
Michalkiewicz J *see* Chrzanowska KH *et al*
Michie S *et al*. Predictive genetic testing in children and adults: a study of emotional impact, **519**
Mieli-Vergani G *see* Kelly AL *et al*
Mignon-Ravix C *et al*. Maternally inherited duplication of the possible imprinted 14q31 region: *letter*, **343**
Mikelsaar RV *et al*. Mosaic terminal del(19)(q13.33:) in a girl with seizures and mental retardation: *letter*, **e2**
Miners AH *see* MacDermot KD *et al*
see MacDermot KD *et al*
Miosczka H *see* Bender BU *et al*
Mischung C *see* Luijten M *et al*
Missirian C *see* Giraudeau F *et al*
Mitchell EA *see* Liu HX *et al*
Moerman P *see* Witters I *et al*
Mogensen J *see* Andersen PS *et al*
Mogensen J *et al*. Development and application of linkage analysis in genetic diagnosis of familial hypertrophic cardiomyopathy: *letter*, **193**
Momoi T *see* Yorifuji T *et al*
Monk D *et al*. No evidence for mosaicism in Silver-Russell syndrome: *letter*, **e11**
Monnier N *et al*. A novel 3600+11.5 kb C>G homozygous splicing mutation in a black African, consanguineous CF family: *letter*, **e4**
Montera M *et al*. A silent mutation in exon 14 of the *APC* gene is associated with exon skipping in a FAP family : *letter*, **863**
Moolman-Smook JC *see* Andersen PS *et al*
Moore AT *see* Payne AM *et al*
Moore GE *see* Hitchins MP *et al*
see Monk D *et al*
Morell RJ *see* Ben-Yosef T *et al*
Moreno F *see* Moreno-Pelayo MA *et al*
Moreno-Pelayo MA *et al*. A cysteine substitution in the zona pellucida domain of α-tectorin results in autosomal dominant, postlingual, progressive, mid frequency hearing loss in a Spanish family: *letter*, **e13**
Morgan G *see* Jones KJ *et al*
see Kwok JBJ *et al*
Morichon N *see* Amiel J *et al*
Morra E *see* Pescarmona GP *et al*
Morreau H *see* Wagner A *et al*
Morris AG *see* Payne AM *et al*
Morris NH *see* O'Shaughnessy KM *et al*
Morrison N *see* Tobias S *et al*
Morrow BE *see* Dar P *et al*
Mortemousque B *see* Coupry I *et al*
Mouridsen HT *see* Bergthorsson JT *et al*
Mubaidin A *see* Hampshire DJ *et al*
Mueller RF *see* Houseman MJ *et al*
see Hutchin TP *et al*
see Kelsell DP *et al*
see van Laer L *et al*
Mugneret F *see* Mignon-Ravix C *et al*
Muller T *see* Haberlandt E *et al*
see Janecke AR *et al*
Muller-Höcker J *see* Jaksch M *et al*
Munnich A *see* Amiel J *et al*
see Bahuau M *et al*
see de Leersnyder H *et al*
see de Lonlay P *et al*
see Edery P *et al*

see Faivre L *et al*
see Francannet C *et al*
see Huber C *et al*
see Huber C *et al*
see Imessaoudene B *et al*
see Quintana-Murci L *et al*
see Saugier-Veber P *et al*
see von Kleist-Retzow J-C *et al*
Mura C *et al.* Variation of iron loading expression in C282Y homozygous haemochromatosis probands and sib pairs: *letter*, **632**
Murday V *see* Barker K *et al*
Muroi J *see* Yorifuji T *et al*
Muroya K *et al.* GATA3 abnormalities and the phenotypic spectrum of HDR syndrome, **374**
Murray A *see* Syrris P *et al*
Musebeck J *see* Tönnies H *et al*
Mustapha M *et al.* Autosomal recessive non-syndromic hearing loss in the Lebanese population: prevalence of the 30delG mutation and report of two novel mutations in the connexin 26 (*GJB2*) gene, **e36**
Mutirangura A *see* Ungaro P *et al*
Myllyluoma B *see* Nopola-Hemmi J *et al*
Myring J *see* Fokstuen S *et al*
see Fokstuen S *et al*

Nadal M *et al.* Clinical and cytogenetic characterisation of a patient with Down syndrome resulting from a 21q22.1→qter duplication: *letter*, **73**
Nagai T *see* Muroya K *et al*
Nagamatsu K *see* Yabe I *et al*
Nakahata T *see* Yorifuji T *et al*
Nakahori Y *see* Kanayama H-O *et al*
Nanba E *see* Yamamoto T *et al*
Nance WE *see* Ben-Yosef T *et al*
Nanko S *see* Ross NLJ *et al*
Naroda T *see* Kanayama H-O *et al*
Nathanson KL *see* Eng C *et al*
Navas C *see* Moreno-Pelayo MA *et al*
Naz S *see* Ben-Yosef T *et al*
Neitzel H *see* Tönnies H *et al*
Nelson CC *see* Erickson RP *et al*
Nelson R *see* Scoggan KA *et al*
Neri G *see* Rossi E *et al*
Neu N *see* Janecke AR *et al*
Neumann HPH *see* Bender BU *et al*
Neumann L *see* Tönnies H *et al*
Neumann LM *see* Tönnies H *et al*
Newton V *see* Hutchin TP *et al*
Nicholls AC *et al.* Homozygosity for a splice site mutation of the *COL1A2* gene yields a non-functional proα2(I) chain and an EDS/OI clinical phenotype: *letter*, **132**
Nicholls RD *see* Lossie AC *et al*
Niebuhr A *see* Bergthorsson JT *et al*
Niebuhr E *see* Bergthorsson JT *et al*
Nielsen KV *see* Bergthorsson JT *et al*
Niemann TH *see* Chadwick RB *et al*
Nieminen P *see* Pirinen S *et al*
Niermeijer M *see* Maat-Kievit A *et al*
Niermeijer MF *see* de Vries BBA *et al*
see Duisterhof M *et al*
see Wagner A *et al*
Ninomiya H *see* Yamamoto T *et al*
Nishimura G *see* Ogata T *et al*
Nishizawa M *see* Yabe I *et al*
Nitowsky HM *see* Dar P *et al*
Nogueira P *see* Amaral MD *et al*
Nolet S *see* Kasprzak L *et al*
Nopola-Hemmi J *et al.* A dominant gene for developmental dyslexia on chromosome 3, **658**
Nord B *see* Kytölä S *et al*
North KN *see* Jones KJ *et al*
Northrup H *see* Khare L *et al*
Notarangelo LD *see* Clementi R *et al*
Nothwang H-G *see* Borck G *et al*
Nur-E-Kamal M *see* Bayoumi R *et al*
Nurnberg G *see* Bayoumi R *et al*
Nurnberg P *see* Luijten M *et al*

Ochs HD *see* Kobayashi I *et al*
Ockenga J *et al.* Low prevalence of *SPINK1* gene mutations in adult patients with chronic idiopathic pancreatitis: *letter*, **243**
Oda T *see* Yabe I *et al*
Odent S *see* Huber C *et al*
Oei PTSP *see* Liu HX *et al*
Offiah AC *et al.* Sponastrime dysplasia: presentation in infancy: *letter*, **889**
Ogata T *see* Boucher CA *et al*
see Muroya K *et al*
Ogata T *et al. SHOX* haploinsufficiency and overdosage: impact of gonadal function status: *review*, **1**
Oguzkaya F *see* Dundar M *et al*
Ohnishi A *see* Yabe I *et al*
Ohno K *see* Sahashi K *et al*
Okano M *see* Kobayashi I *et al*
Olafsdóttir G *see* Hrafnkelsson J *et al*
Oliver E *see* Vincent A *et al*
Ollikainen V *see* Nopola-Hemmi J *et al*
Olopade OI *see* Fackenthal JD *et al*
Olschewski M *see* Bender BU *et al*
Olschwang S *et al.* Peutz-Jeghers families unlinked to *STK11/LKB1* gene mutations are highly predisposed to primitive biliary adenocarcinoma, **356**
Olsen JH *see* Bergthorsson JT *et al*
Olveira C *see* Amaral MD *et al*
Orfanos CE *see* Zouboulis CC *et al*
Orii T *see* Imamura A *et al*
Oriola J *see* Kasprzak L *et al*
Orszagh M *see* Bender BU *et al*
Osborn AH *see* Houseman MJ *et al*
O'Shaughnessy KM *et al.* Thrombophilic polymorphisms in pre-eclampsia: altered frequency of the functional 98C>T polymorphism of glycoprotein IIIa: *letter*, **775**
Östman-Smith I *see* Blair E *et al*
Ostrander E *see* Eng C *et al*
Osumi T *see* Imamura A *et al*
Ouvrier RA *see* Jones KJ *et al*
Overhauser J *see* Cerruti Mainardi P *et al*
Ozkul Y *see* Dundar M *et al*
Ozyazgan I *see* Dundar M *et al*

Pace JM *et al.* Disruption of one intra-chain disulphide bond in the carboxyl-terminal propeptide of the proα1(I) chain of type I procollagen permits slow assembly and secretion of overmodified, but stable procollagen trimers and results in mild osteogenesis imperfecta, **443**
Pacheco P *see* Amaral MD *et al*
Padovani P *see* Huber C *et al*
Page DC *see* Jaruzelska J *et al*
Pages-Berhouet S *see* Gad S *et al*
Pagès S *see* Chompret A *et al*
Pagnamenta A *see* Houseman MJ *et al*
Pandya A *see* Ben-Yosef T *et al*
Pang C-P *see* Lam C-W *et al*
Pannier E *see* Viot G *et al*
Parenti G *see* Filocamo M *et al*
Parent P *see* Giraudeau F *et al*
Parisi MA *et al.* The spectrum and evolution of phenotypic findings in *PTEN* mutation positive cases of Bannayan-Riley-Ruvalcaba syndrome: *letter*, **52**
Parker M *see* Hutchin TP *et al*
Parsons JM *see* Henwood J *et al*
Paskaleva I *see* Horvath A *et al*
Passarge E *see* Albrecht B *et al*
Pastore G *see* Cerruti Mainardi P *et al*
Paterson AD *see* Dar P *et al*
Patsch JR *see* Eller P *et al*
see Clementi R *et al*
Pavia C *see* Kasprzak L *et al*
Pawelczyk L *see* Jaruzelska J *et al*
Payne AM *et al.* Clustering and frequency of mutations in the retinal guanylate cyclase (*GUCY2D*) gene in patients with dominant cone-rod dystrophies: *letter*, **611**
Pearson P *see* van Tilburg J *et al*
Pedespan D *see* de Lonlay P *et al*
Pelet A *see* Bahuau M *et al*
Pellestor F *see* Lefort G *et al*
Peltomäki P *see* Liu T *et al*
Penet C *see* Huber C *et al*
Penque D *see* Amaral MD *et al*
Pérez-Frías J *see* Amaral MD *et al*
Perfumo C *see* Cerruti Mainardi P *et al*
see Rossi E *et al*
Perrin-Schmitt F *see* Dollfus H *et al*
Pescarmona GP *et al.* Haptoglobin genotype as a risk factor for postmenopausal osteoporosis: *letter*, **636**
Peters H *see* Hegde MR *et al*
Peters WHM *see* Zusterzeel PLM *et al*
Petit C *see* Mustapha M *et al*
Philippe C *see* Bourdon V *et al*
Piaggio F *see* Montera M *et al*
Piccini F *see* Rossi E *et al*
Pickard C *see* Henwood J *et al*
Pierluigi M *see* Cerruti Mainardi P *et al*
see Rossi E *et al*
Pignatti P *see* Castellani C *et al*
Pigott RW *see* Turnpenny PD and Pigott RW
Pin I *see* Monnier N *et al*
Pingault V *et al.* A heterozygous endothelin 3 mutation in Waardenburg-Hirschsprung disease: is there a dosage effect of *EDN3/EDNRB* gene mutations on neurocristopathy phenotypes?: *letter*, **205**
Pinto M *see* Gad S *et al*

Pirinen S *et al.* Recessively inherited lower incisor hypodontia: *letter*, **551**
Pirinen S *see* Hannula K *et al*
Plass C *see* Costello JF and Plass C
Polak M *see* Faivre L *et al*
Pongratz D *see* Jaksch M *et al*
Pope FM *see* Nicholls AC *et al*
Porter TR *see* Verma L *et al*
Pouliquen JC *see* Faivre L *et al*
Powis SH *see* Veal CD *et al*
Prasad SD *see* van Laer L *et al*
Preece M *see* Monk D *et al*
Preece MA *see* Hitchins MP *et al*
Price SJ *see* Blair E *et al*
Prieur M *see* Mégarbané A *et al*
Priolo M *see* Cinti R *et al*
Priolo M, Laganà C. Ectodermal dysplasias: a new clinical-genetic classification: *review*, **579**
Prior TW *see* Chadwick RB *et al*
Priston M *see* Vincent A *et al*
Pritchard M *see* Nadal M *et al*
Pyatt RE *see* Chadwick RB *et al*

Quan F *see* Tahmaz FE *et al*
Quarrell OWJ *see* de Vries BBA *et al*
see Shannon NL *et al*
Quillet R *see* Dollfus H *et al*
Quintana-Murci L *et al.* Mitochondrial DNA inheritance in patients with deleted mtDNA: *letter*, **e28**

Raclin V *see* Amiel J *et al*
Radhakrishna U *see* Ben-Yosef T *et al*
Raguenes O *see* Mura C *et al*
Ragusa A *see* Borgione E *et al*
see Rossi E *et al*
Rai M *see* Mustapha M *et al*
Rainier S *et al.* Hereditary spastic paraplegia linked to chromosome 14q11-q21: reduction of the SPG3 locus interval from 5.3 to 2.7 cM: *letter*, **e39**
Rajpar MH *see* Verma L *et al*
Rajput K *see* Rickard S *et al*
Ramsden S *see* Watson P *et al*
Ranke MB *see* Eggermann T *et al*
Ranta R *see* Wong FK *et al*
Raskin S *see* Kwok JBJ *et al*
Ratner N *see* Steigerwald K *et al*
Rautio J *see* Wong FK *et al*
Ravazzolo R *see* Cinti R *et al*
Ravine D *see* Fokstuen S *et al*
Ray PN *see* Grunebaum E *et al*
Reardon W *see* Houseman MJ *et al*
see Kelsell DP *et al*
Reardon W *et al.* A novel germline mutation of the *PTEN* gene in a patient with macrocephaly, ventricular dilatation, and features of VATER association, **820**
Rees DRZ *see* Baker L *et al*
Rees MI *see* Baker L *et al*
Regan R *see* de Vries BBA *et al*
Regis S *see* Filocamo M *et al*
Reid E *et al.* The spastic paraplegia SPG10 locus: narrowing of critical region and exclusion of sodium channel gene *SCN8A* as a candidate: *letter*, **65**
Reis A *see* Bayoumi R *et al*
see Chrzanowska KH *et al*
see Maraschio P *et al*
see Vanita *et al*
Renvert S *see* Zhang Y *et al*
Resta N *see* Montera M *et al*
Reuser AJJ *see* Ausems MGEM *et al*
Reynolds J *see* Ungaro P *et al*
Riazuddin S *see* Ben-Yosef T *et al*
see Ben-Yosef T *et al*
Richards FM *see* Verma L *et al*
Richards M *see* Hallowell N *et al*
Richards SK *see* Chadwick RB *et al*
Richardson A-L *see* Fackenthal JD *et al*
Rickard S *see* Houseman MJ *et al*
Rickard S *et al.* Recurrent mutations in the deafness gene *GJB2* (connexin 26) in British Asian families: *letter*, **530**
Ricordy R *see* Maraschio P *et al*
Riegel M *see* Kotzot D *et al*
Rigby AS *see* Shannon NL *et al*
Rihkanen H *see* Aaltonen L-M *et al*
Rip J *see* Coulter-Mackie MB *et al*
Rivera-Fillat F *see* Kasprzak L *et al*
Road F *see* Hallowell N *et al*
Roberts E *see* Hampshire DJ *et al*
see Henwood J *et al*
Robertson NJ *see* Boardman JP *et al*
Robinson BG *see* Fackenthal JD *et al*
Rocha J *see* Seixas S *et al*
Rodeck CH *see* Kalache KD *et al*
Rodrigues F *see* Kelly AL *et al*
Roelofs HMJ *see* Zusterzeel PLM *et al*
Roggenbuck J *see* Ehrlich M *et al*
Roifman CM *see* Grunebaum E *et al*
Romana SP *see* Mégarbané A *et al*
Romano C *see* Borgione E *et al*
see Rossi E *et al*
Romero L *see* Moreno-Pelayo MA *et al*
Roos RAC *see* Duisterhof M *et al*
Ropers H-H *see* Borck G *et al*
Rose H *see* Kirchhoff M *et al*
Ross F *see* Eccles D *et al*
Ross NLJ *et al.* Triplication of several PAR1 genes and part of the *Homo sapiens* specific Yp11. 2/Xq21. 3 region of homology in a 46,X,t(X;Y)(p22. 33;p11. 2) male with schizophrenia: *letter*, **710**
Rossi E *et al.* Cryptic telomeric rearrangements in subjects with mental retardation associated with dysmorphism and congenital malformations: *letter*, **417**
Roth S *see* Aaltonen L-M *et al*
Röthlisberger B *see* Kotzot D *et al*
Röthlisberger B *et al.* Supernumerary marker chromosome (1) of paternal origin and maternal uniparental disomy 1 in a developmentally delayed child : *letter*, **885**
Rötig A *see* Quintana-Murci L *et al*
see von Kleist-Retzow J-C *et al*
Roubertie A *see* Tuffery-Giraud S *et al*
Roumelioti KN *see* Ausems MGEM *et al*
Rowland J *see* van Laer L *et al*
Rowland JS *et al.* A comparison of methods for gene dosage analysis in HMSN type 1, **90**
Royer G *see* Imessaoudene B *et al*
Roylance R *see* Tomlinson IPM *et al*
Rubinsztein DC *see* Ho LW *et al*
see McConnell RS *et al*
see Reid E *et al*
see Taylor AE *et al*
Ruiz-Perez VL *see* Tompson SWJ *et al*
Ruiz-Ponte C *et al.* The Asp1822Val variant of the *APC* gene is a common polymorphism without clinical implications: *letter*, **e33**
Rumeur S *see* Huber C *et al*
Rupar CA *et al.* A G339R mutation in the *CTNS* gene is a common cause of nephropathic cystinosis in the south western Ontario Amish Mennonite population: *letter*, **615**
Rupar C *see* Rupar CA *et al*
Rupprecht E *see* Horn D *et al*
Russell K *see* Faivre L *et al*
Russo S *see* Monk D *et al*
Rustin P *see* Quintana-Murci L *et al*
see von Kleist-Retzow J-C *et al*

Saar K *see* Bayoumi R *et al*
Saha N *see* Liu Y *et al*
Sahashi K *see* Sahashi K *et al*
Sahashi K *et al.* Functional characterisation of mitochondrial $tRNA^{Tyr}$ mutation (5877G→A) associated with familial chronic progressive external ophthalmoplegia, **703**
Saheki T *see* Hwu W-L *et al*
Sakiyama Y *see* Kobayashi I *et al*
Salahshor S *see* Liu T *et al*
Salazar-De-Sousa J *see* Seixas S *et al*
Salem N *see* Mustapha M *et al*
Salgueiro C *see* Seixas S *et al*
Salmela PI *see* Kytölä S *et al*
Salovaara R *see* Avizienyte E *et al*
Sam S *see* Tahmaz FE *et al*
Samaras L *see* Mégarbané A *et al*
Sancandi M *see* Pingault V *et al*
Sanders L *see* Mahavni V *et al*
Sandkuijl LA *see* Ausems MGEM *et al*
Santoro IM *see* Steigerwald K *et al*
Santra A *see* Kadakol A *et al*
Sanz R *see* Lorda-Sanchez I *et al*
Sappal BS *see* Kadakol A *et al*
Sarda P *see* de Lonlay P *et al*
see Lefort G *et al*
Sargent CA *see* Boucher CA *et al*
see Ross NLJ *et al*
Sarhadi V *see* Vanita *et al*
Sasaki H *see* Komeichi K *et al*
see Yabe I *et al*
Saudubray JM *see* de Lonlay P *et al*
Saugier-Veber P *et al.* Detection of heterozygous *SMN1* deletions in SMA families using a simple fluorescent multiplex PCR method: *letter*, **240**
Sauper T *see* Eller P *et al*
Savoldi G *see* Clementi R *et al*
Savov A *see* Horvath A *et al*
Scharfe C *see* Jaksch M *et al*
Scheuner MT *see* Gad S *et al*
Schinzel A *see* Kotzot D *et al*
see Mikelsaar RV *et al*
see Röthlisberger B *et al*

Schmidt A *see* Fritz B *et al*
Schmidt D *see* Bender BU *et al*
Schmitz J *see* de Lonlay P *et al*
Schneider M *see* Clementi R *et al*
Schneider T *see* Kirches E *et al*
Schofield PR *see* Kwok JBJ *et al*
Schteingart D *see* Stratakis CA *et al*
Schulenberg E *see* Talks SJ *et al*
Schulze I *see* Tönnies H *et al*
Schwartz CE *see* Villard L *et al*
Schwerdtfeger R *see* Fritz B *et al*
Schwerdtfeger UF *see* Fritz B *et al*
Scoggan KA *et al.* Identification of two novel mutations in the *CACNA1A* gene responsible for episodic ataxia type 2: *letter*, **249**
Scotet V *see* Mura C *et al*
Scott HS *see* Ben-Yosef T *et al*
Scott RJ *see* Kalache KD *et al*
Seishu S *see* Muroya K *et al*
Seixas S *et al.* Association between the defective Pro369Ser mutation and in vivo intrahepatic α1-antitrypsin accumulation: *letter*, **472**
Serdaroglu P *see* Akbas F *et al*
Seri M *see* Cinti R *et al*
Seta N *see* de Lonlay P *et al*
Sham MH *et al.* Novel mutations of *SOX10* suggest a dominant negative role in Waardenburg-Shah syndrome: *letter*, **e30**
Shannon NL *et al.* An epidemiological study of Wolf-Hirschhorn syndrome: life expectancy and cause of mortality, **674**
Shenton A *see* Hopwood P *et al*
Sheridan E *see* Flintoff KJ *et al*
Shiari R *see* Kobayashi I *et al*
Shibuya K *see* Ben-Yosef T *et al*
Shimozawa N *see* Imamura A *et al*
Shipley J *see* Barker K *et al*
Short MP *see* Blakley P *et al*
Shoubridge EA *see* von Kleist-Retzow J-C *et al*
Sidi D *see* de Leersnyder H *et al*
Siffroi JP *et al.* Alternative centromeric inactivation in a pseudodicentric t(Y;13)(q12;p11. 2) translocation chromosome associated with extreme oligozoospermia: *letter*, **802**
Sigvaldason H *see* Hrafnkelsson J *et al*
Silengo M *see* Cinti R *et al*
Silva L *see* Seixas S *et al*
Simon DK *et al.* A frameshift mitochondrial complex I gene mutation in a patient with dystonia and cataracts: is the mutation pathogenic?: *letter*, **58**
Simpson SA *et al.* Prenatal testing for Huntington's disease: experience within the UK 1994-1998: *letter*, **333**
Singh D *see* Vanita *et al*
Singh JR *see* Vanita *et al*
Sirimana T *see* Rickard S *et al*
Siwarski D *et al.* Identification of a transcriptionally compromised allele of *c-MYC* in a North American family: *letter*, **47**
Sklansky M *see* Vitale E *et al*
Smahi A *see* Siffroi JP *et al*
Smalley SL *see* Khare L *et al*
Smeets D *see* Hiel JAP *et al*
Smith M *see* Khare L *et al*
Smith RJH *see* Fransen E *et al*
see Haberlandt E *et al*
see van Laer L *et al*
Smithson SF *see* de Vries BBA *et al*
Soininen H *see* Finnilä S *et al*
Soriano C *see* Coupry I *et al*
Souraty N *see* Mégarbané A *et al*
Sperling K *see* Vanita *et al*
Splitt M *see* de Vries BBA *et al*
Spranger J *see* Horn D *et al*
Zur Stadt U *see* Clementi R *et al*
Stallings RL *see* McEntagart M *et al*
Stanier P *see* Hitchins MP *et al*
see Monk D *et al*
Stavropoulou C *see* Mignon-Ravix C *et al*
StClair D *see* Lambert J-C *et al*
Steegers EAP *see* Zusterzeel PLM *et al*
Steel Z *see* Meiser B *et al*
Steering Committee of the Breast Cancer Information Consortium Cove (BIC) *see* Eng C *et al*
Steffensen U *see* Mogensen J *et al*
Steigerwald K *et al.* A distinct splice form of *APC* is highly expressed in neurones but not commonly mutated in neuroepithelial tumours: *letter*, **257**
Stella A *see* Montera M *et al*
Stevens MW *see* Chadwick RB *et al*
Stöckl B *see* Haberlandt E *et al*
Stoetzel C *see* Dollfus H *et al*
Stoppa-Lyonnet D *see* Bougeard G *et al*
see Chompret A *et al*
see Gad S *et al*
Stratakis CA *see* Zouboulis CC *et al*
Stratakis CA *et al.* A syndrome of overgrowth and acromegaloidism with normal growth hormone secretion is associated with chromosome 11 pericentric inversion: *letter*, **338**
Segmental uniparental isodisomy (UPD) for 2p16 without clinical symptoms: implications for UPD and other genetic studies of chromosome 2: *letter*, **106**
Strizheva GD *see* Khare L *et al*
Stroppiano M *see* Filocamo M *et al*
Stuhrmann M *see* Hughes D *et al*
see Ockenga J *et al*
Stumm M *see* Chrzanowska KH *et al*
see Tönnies H *et al*
Sulmont V *see* Faivre L *et al*
see Huber C *et al*
Sumption ND, Barber JCK. A transmitted deletion of 2q13 to 2q14.1 causes no phenotypic abnormalities: *letter*, **125**
Super M *see* de Vries BBA *et al*
see Watson P *et al*
Surmacz T *see* Jaruzelska J *et al*
Surry S *see* Rupar CA *et al*
Sutherland J *see* Vincent A *et al*
Suvari A *see* Mikelsaar RV *et al*
Suzuki Y *see* Imamura A *et al*
see Yabe I *et al*
Sybert VP *see* Parisi MA *et al*
Syrris P *see* Boardman JP *et al*
Syrris P *et al.* Mutation detection in long QT syndrome: a comprehensive set of primers and PCR conditions, **705**
Syrrou M, Fryns J-P. Interstitial deletion of chromosome 11 (q22.3-q23.2) in a boy with mild developmental delay: *letter*, **621**
Szabo C *see* Eng C *et al*

Taanman J-W *see* von Kleist-Retzow J-C *et al*
Tahmaz FE *et al.* A partial deletion of the aspartoacylase gene is the cause of Canavan disease in a family from Mexico: *letter*, **e9**
Taine L *see* Coupry I *et al*
see Giraudeau F *et al*
Taipale M *see* Nopola-Hemmi J *et al*
Takahashi M *see* Kanayama H-O *et al*
Takei A *see* Yabe I *et al*
Takiyama Y *see* Yabe I *et al*
Talks SJ *et al.* De novo mutations in the 5' regulatory region of the Norrie disease gene in retinopathy of prematurity, **e46**
Tam PKH *see* Sham MH *et al*
Tamanini A *see* Castellani C *et al*
Tanaka M *see* Sahashi K *et al*
Tantau J *see* Viot G *et al*
Tanzarella C *see* Maraschio P *et al*
Tardieu M *see* Bourdon V *et al*
see Villard L *et al*
Tarnopolsky MA *see* Simon DK *et al*
Tarttelin EE *et al.* Molecular genetic heterogeneity in autosomal dominant drusen, **381**
Tashiro K *see* Komeichi K *et al*
see Yabe I *et al*
Tassone F *et al.* A majority of fragile X males with methylated, full mutation alleles have significant levels of *FMR1* messenger RNA, **453**
Tatakis DN *see* Zhang Y *et al*
Tavtigian SV *see* Eng C *et al*
Taylor AE *et al.* Genetic association of an *LBP-1c/CP2/LSF* gene polymorphism with late onset Alzheimer's disease, **232**
Taylor AK *see* Tassone F *et al*
Taylor GR *see* Flintoff KJ *et al*
see Houseman MJ *et al*
see Rowland JS *et al*
see van Laer L *et al*
Taymans SE *see* Stratakis CA *et al*
Ten Berg T *see* Ausems MGEM *et al*
Teh BT *see* Kanayama H-O *et al*
see Kytölä S *et al*
see Wong FK *et al*
Ten Kate LP *see* Henneman L *et al*
Tenconi R *see* Rossi E *et al*
Theriaque D *see* Lossie AC *et al*
Thesleff I *see* Pirinen S *et al*
Thomas G *see* Olschwang S *et al*
Thompson KR *see* Hutchin TP *et al*
Thompson PW *see* Baker L *et al*
Thomson JDR *see* Henwood J *et al*
Tibben A *see* Duisterhof M *et al*
see Maat-Kievit A *et al*
Tiberio G *see* Giannotti A *et al*
Tiepolo L *see* Maraschio P *et al*
Tillman D *see* Veal CD *et al*
Tobias S *et al.* Absence of learning difficulties in a hyperactive boy with a terminal Xp deletion encompassing the *MRX49* locus: *letter*, **466**
Tobias V *see* Jones KJ *et al*
Tolmie JL *see* Dundar M *et al*
see Tobias S *et al*

Tomlinson IPM *et al.* Two hits revisited again: *review*, **81**
Tommerup N *see* Borck G *et al*
Tompson SWJ *et al.* Ellis-van Creveld syndrome resulting from segmental uniparental disomy of chromosome 4: *letter*, **e18**
Tönnies H *see* Borck G *et al*
Tönnies H *et al.* De novo terminal deletion of chromosome 15q26.1 characterised by comparative genomic hybridisation and FISH with locus specific probes: *letter*, **617**
Two further cases of WHS with unbalanced de novo translocation t(4;8) characterised by CGH and FISH: *letter*, **e21**
Tops C *see* Wagner A *et al*
Toro JR *see* Stratakis CA *et al*
Toutain A *see* Giraudeau F *et al*
Trainer AH *see* de Vries BBA *et al*
Trembath RC *see* Veal CD *et al*
Trijsburg RW *see* Duisterhof M *et al*
Trump D *see* Willatt L *et al*
Trzepacz C *see* Steigerwald K *et al*
Tsang M-W *see* Lam C-W *et al*
Tsien F *see* Ehrlich M *et al*
Tsukamoto T *see* Imamura A *et al*
Tuck-Muller CM *see* Ehrlich M *et al*
Tucker K *see* Meiser B *et al*
Tuffery-Giraud S *et al.* No evidence of allelic heterogeneity in the *DYT1* gene of European patients with early onset torsion dystonia: *letter*, **e35**
Tulinius H *see* Hrafnkelsson J *et al*
Tummler B *see* Amaral MD *et al*
Turleau C *see* Mégarbané A *et al*
Turner G *see* Flintoff KJ *et al*
Turner ML *see* Stratakis CA *et al*
Turnpenny PD, Pigott RW. Deletion 22q11 syndrome: acknowledging a lost eponym as we say farewell to an acronym: *letter*, **271**
Tzetis M *see* Amaral MD *et al*

Uematsu A *see* Yorifuji T *et al*
Umeå S8 *see* Liu T *et al*
Ungaro P *et al.* Molecular characterisation of four cases of intrachromosomal triplication of chromosome 15q11-q14, **26**
Unsinn K *see* Janecke AR *et al*
Upadhyaya M *see* Baker L *et al*
Usa *see* Martin DM *et al*
Utermann G *see* Haberlandt E *et al*
see Janecke AR *et al*
Uutela A *see* Aktan-Collan K *et al*
Uygur C *see* Zhang Y *et al*

Vaheri A *see* Aaltonen L-M *et al*
Välimäki M *see* Kytölä S *et al*
Valler D *see* Nicholls AC *et al*
van Assche A *see* Witters I *et al*
van Beersum SEC *see* Bongers EMHF *et al*
van Bokhoven H *see* Bongers EMHF *et al*
van Camp G *see* Fransen E *et al*
see van Laer L *et al*
van de Heyning CM *see* van Laer L *et al*
van de Heyning PH *see* Fransen E *et al*
van den Boer-van den Berg HM, Maat-Kievit AA. The whole truth and nothing but the truth, but what is the truth?, **39**
van den Boer-van den Berg H *see* Maat-Kievit A *et al*
van den Heuvel LPWJ *see* Hiel JAP *et al*
van der Burgt I *see* Hiel JAP *et al*
van der Horst HE *see* Henneman L *et al*
van der Luijt RB *et al.* De novo recurrent germline mutation of the *BRCA2* gene in a patient with early onset breast cancer, **102**
van der Meer C *see* Wagner A *et al*
van der Ploeg HM *see* Henneman L *et al*
van der Sijs-Bos CJM *see* van der Luijt RB *et al*
van Engelen BGM *see* Hiel JAP *et al*
van Haeften TW *see* van Tilburg J *et al*
Vanita *et al.* A unique form of autosomal dominant cataract explained by gene conversion between β-crystallin B2 and its pseudogene: *letter*, **392**
van Kien PK *see* Mignon-Ravix C *et al*
van Krimpen C *see* Wagner A *et al*
van Laer L *et al.* A common founder for the 35delG *GJB2* gene mutation in connexin 26 hearing impairment, **515**
van Maldergem L *see* Fransen E *et al*
van Ommen G-J *see* Maat-Kievit A *et al*
van Schaftingen E *see* de Lonlay P *et al*
van Schoubroeck D *see* Witters I *et al*
van Thienen M-N *see* Bongers EMHF *et al*
van Tilburg J *et al.* Defining the genetic contribution of type 2 diabetes mellitus: *review*, **569**
van Velthoven V *see* Bender BU *et al*
van Zon PHA *see* van der Luijt RB *et al*
Varb K *see* Mikelsaar RV *et al*
Varesco L *see* Montera M *et al*
see Steigerwald K *et al*
Varilo T *see* Pirinen S *et al*
Varon R *see* Maraschio P *et al*
see Chrzanowska KH *et al*
Varotto S *see* Clementi R *et al*
Vasen HFA *see* Wagner A *et al*
Vásquez C *see* Amaral MD *et al*
Veal CD *et al.* Identification of a novel psoriasis susceptibility locus at 1p and evidence of epistasis between *PSORS1* and candidate loci, **7**
Vega A *see* Ruiz-Ponte C *et al*
Vekemans M *see* de Leersnyder H *et al*
see Mégarbané A *et al*
Vergnaud G *see* Giraudeau F *et al*
Verhagen WIM *see* Fransen E *et al*
Verlingue C *see* Amaral MD *et al*
Verma L *et al.* Germline mutation analysis of the transforming growth factor β receptor type II (*TGFBR2*) and E-cadherin (*CDH1*) genes in early onset and familial colorectal cancer: *letter*, **e7**
Verstreken M *see* Fransen E *et al*
Vial E *see* Saugier-Veber P *et al*
Vidaud D *see* Bahuau M *et al*
Vidaud M *see* Bahuau M *et al*
Vieira I *see* Amaral MD *et al*
Vierimaa O *see* Kytölä S *et al*
Vigo CG *see* Nadal M *et al*
Vijg J *see* Eng C *et al*
Villablanca A *see* Kytölä S *et al*
Villabona C *see* Kasprzak L *et al*
Villain E *see* de Leersnyder H *et al*
Villamar M *see* Moreno-Pelayo MA *et al*
Villard L *et al.* Segregation of a totally skewed pattern of X chromosome inactivation in four familial cases of Rett syndrome without *MECP2* mutation: implications for the disease, **435**
Villeneuve N *see* de Lonlay P *et al*
Vincent A *et al.* Phenotypic heterogeneity of *CYP1B1*: mutations in a patient with Peters' anomaly: *letter*, **324**
Viot G *et al.* A new case of exomphalos, short limbs, and macrogonadism syndrome: *letter*, **e8**
Virgilii F *see* Giannotti A *et al*
Visser W *see* Zusterzeel PLM *et al*
Vitale E *et al.* Suggestive linkage of situs inversus and other left-right axis anomalies to chromosome 6p: *letter*, **182**
Volleth M *see* Tönnies H *et al*
von Kleist-Retzow J-C *et al.* Mutations in *SURF1* are not specifically associated with Leigh syndrome: *letter*, **109**
Voutilainen A *see* Nopola-Hemmi J *et al*
Vuust J *see* Andersen PS *et al*

Wadekar R *see* Ross NLJ *et al*
Wagner A *et al.* Atypical HNPCC owing to *MSH6* germline mutations: analysis of a large Dutch pedigree, **318**
Wagner TMU *see* Eng C *et al*
Wallis S *see* Nicholls AC *et al*
Walter MA *see* Vincent A *et al*
Wandall A *see* Fritz B *et al*
Wang J-H *see* Hwu W-L *et al*
Wang R *see* Barker K *et al*
Warich-Kirches M *see* Kirches E *et al*
Wárlám-Rodenhuis CC *see* van der Luijt RB *et al*
Warren ST *see* Hegde MR *et al*
Watkins H *see* Blair E *et al*
Watson P *et al.* Angelman syndrome phenotype associated with mutations in *MECP2*, a gene encoding a methyl CpG binding protein, **224**
Wattenhofer M *see* Ben-Yosef T *et al*
Watterson KG *see* Henwood J *et al*
Weatherall D. Cyril Astley Clarke, **281**
Webb T *see* Whittington JE *et al*
Webb T, Latif F. Rett syndrome and the *MECP2* gene: *review*, **217**
Weber J *see* Vitale E *et al*
Weemaes CMR *see* Hiel JAP *et al*
Weis S *see* Kirches E *et al*
Weleber RG *see* Tarttelin EE *et al*
Welkenhuysen M *et al.* A community based study on intentions regarding predictive testing for hereditary breast cancer: *letter*, **540**
Westerveld A *see* Luijten M *et al*
White SM *see* de Vries BBA *et al*
Whitney MM *see* Lossie AC *et al*
Whittington JE *et al.* Population prevalence and estimated birth incidence and mortality rate for people with Prader-Willi syndrome in one UK Health Region: *letter*, **792**
Widén E *see* Nopola-Hemmi J *et al*
Wijmenga C *see* Ausems MGEM *et al*
see van Tilburg J *et al*
Wijnen J *see* Wagner A *et al*
Wilcox ER *see* Ben-Yosef T *et al*
Wilkinson AG *see* Dundar M *et al*
Wilkinson I *see* Jones KJ *et al*
Willatt L *et al.* Satellites on the terminal short arm of chromosome 12 (12ps), inherited through several generations in three families: a new variant without phenotypic effect: *letter*, **723**
Willatt LR *see* Dee SL *et al*

Williams CA *see* Lossie AC *et al*
Williams NA *see* Ross NLJ *et al*
Williams-Lyn D *see* Vincent A *et al*
Williamson R *see* Gaff C *et al*
Willing MC *see* Pace JM *et al*
Winter RM *see* Akrami SM *et al*
see de Vries BBA *et al*
see de Vries BBA *et al*
see Offiah AC *et al*
Winther K *see* Bergthorsson JT *et al*
Wirth J *see* Borck G *et al*
Witte M *see* Erickson RP *et al*
Witters I *et al*. Pregnancy outcome and long term prognosis in 868 children born after second trimester amniocentesis for maternal serum positive triple test screening and normal prenatal karyotype: *letter*, **336**
Wojda A *see* Jaruzelska J *et al*
Wolf JP *see* Siffroi JP *et al*
Wollmann HA *see* Eggermann T *et al*
Wong FK *see* Kanayama H-O *et al*
see Kytölä S *et al*
Wong FK *et al*. Genetic heterogeneity and exclusion of a modifying locus at 17p11.2-p11.1 in Finnish families with van der Woude syndrome: *letter*, **198**
Woods CG *see* Hampshire DJ *et al*
Wriekat A-L *see* Hampshire DJ *et al*
Wright MJ *see* Tompson SWJ *et al*
Wuyts FL *see* Fransen E *et al*
Wyttenbach A *see* Ho LW *et al*

Xiang F *see* Villard L *et al*
Xuereb J *see* Taylor AE *et al*

Yabe I *see* Komeichi K *et al*
Yabe I *et al*. Predisposing chromosome for spinocerebellar ataxia type 6 (SCA6) in Japanese: *letter*, **328**
Yamada M *see* Kobayashi I *et al*
Yamaguchi N *see* Hwu W-L *et al*
Yamamoto K *see* Muroya K *et al*
Yamamoto T *et al*. DEFECT 11 syndrome associated with agenesis of the corpus callosum: *letter*, **e5**
Yamanaka C *see* Yorifuji T *et al*
Yamashita I *see* Komeichi K *et al*
see Yabe I *et al*
Yang F *see* Talks SJ *et al*
Yang J *see* Ross NLJ *et al*
Yang T *see* Lam C-W *et al*
Yao J *see* von Kleist-Retzow J-C *et al*
Yara A *see* Kobayashi I *et al*
Yates JRW *see* Dee SL *et al*
Yip A *see* Taylor AE *et al*
Yoneda M *see* Sahashi K *et al*
Yorifuji T *et al*. Analysis of the *SRY* gene in Turner syndrome patients with Y chromosomal material: *letter*, **e41**
Young ID *see* de Vries BBA *et al*
Yu Z *see* Li Y *et al*
Yuksel-Apak M *see* Kayserili H *et al*

Zackai E *see* Faivre L *et al*
Zackai EH *see* Gong W *et al*
Zara F *see* Cerruti Mainardi P *et al*
Zerova TE *see* Röthlisberger B *et al*
Zerres K *see* Albrecht B *et al*
Zhang K *see* Li Y *et al*
Zhang Y *et al*. Evidence of a founder effect for four cathepsin C gene mutations in Papillon-Lefèvre syndrome patients, **96**
Zhang Z *see* Imamura A *et al*
see Villard L *et al*
Zhao X *see* Rainier S *et al*
Zhou X-P *see* Reardon W *et al*
Zollino M *see* Rossi E *et al*
Zori RT *see* Lossie AC *et al*
Zouboulis CC *et al*. Keratosis pilaris/ulerythema ophryogenes and 18p deletion: is it possible that the *LAMA1* gene is involved?: *letter*, 127
Zuffardi O *see* Rossi E *et al*
Zur Stadt U *see* Clementi R *et al*
Zusterzeel PLM *et al*. A polymorphism in the gene for microsomal epoxide hydrolase is associated with pre-eclampsia, **234**
Zwinderman AH *see* Wagner A *et al*

VOLUME 38 • SUBJECT INDEX

11q14 deletion, Leucodystrophy and oculocutaneous albinism in a child with an 11q14 deletion, **35**

14q31 duplication, Maternally inherited duplication of the possible imprinted 14q31 region: *letter*, **343**

15q11-q13, Distinct phenotypes distinguish the molecular classes of Angelman syndrome, **834**

18p deletion, Keratosis pilaris/ulerythema ophryogenes and 18p deletion: is it possible that the *LAMA1* gene is involved?: *letter*, **127**

1p36 deletion syndrome, Use of a set of highly polymorphic minisatellite probes for the identification of cryptic 1p36.3 deletions in a large collection of patients with idiopathic mental retardation: *letter*, **121**

1qter deletions, Submicroscopic subtelomeric 1qter deletions: a recognisable phenotype?: *letter*, **175**

22q11.2 homozygosity, A region of homozygosity within 22q11.2 associated with congenital heart disease: recessive DiGeorge/velocardiofacial syndrome?: *letter*, **533**

3272-26A>G, Cystic fibrosis patients with the 3272-26A>G splicing mutation have milder disease than F508del homozygotes: a large European study: *letter*, **777**

30delG mutation, Autosomal recessive non-syndromic hearing loss in the Lebanese population: prevalence of the 30delG mutation and report of two novel mutations in the connexin 26 (*GJB2*) gene: *letter*, **e36**

35delG, A common founder for the 35delG *GJB2* gene mutation in connexin 26 hearing impairment, **515**

3p22, Identification of a new *TWIST* mutation (7p21) with variable eyelid manifestations supports locus homogeneity of BPES at 3q22: *letter*, **470**

5p deletion, Clinical and molecular characterisation of 80 patients with 5p deletion: genotype-phenotype correlation, **151**

α1-antitrypsin, Association between the defective Pro369Ser mutation and in vivo intrahepatic α1-antitrypsin accumulation: *letter*, **472**

α-tectorin A cysteine substitution in the zona pellucida domain of α-tectorin results in autosomal dominant, postlingual, progressive, mid frequency hearing loss in a Spanish family: *letter*, **e13**

ABCC6, Pseudoxanthoma elasticum: evidence for the existence of a pseudogene highly homologous to the *ABCC6* gene: *letter*, **457**

absent teeth, Distal arthrogryposis: a new type with distinct facial appearance and absent teeth: *letter*, **e22**

acromicric dysplasia, Acromicric dysplasia: long term outcome and evidence of autosomal dominant inheritance, **745**

acyl-CoA oxidase, Temperature sensitive acyl-CoA oxidase import in group A peroxisome biogenesis disorders: *letter*, **871**

adducted thumbs, Adducted thumb-club foot syndrome in sibs of a consanguineous Austrian family: *letter*, **265**

adenomatous polyposis coli, A distinct splice form of *APC* is highly expressed in neurones but not commonly mutated in neuroepithelial tumours: *letter*, **257**

adult onset, A Chinese adult onset type II citrullinaemia patient with 851del4/1638ins23 mutations in the *SLC25A13* gene: *letter*, **e23**

aganglionic megacolon, Hirschsprung disease, associated syndromes, and genetics: a review: *review*, **729**

agenesis of the corpus callosum, DEFECT 11 syndrome associated with agenesis of the corpus callosum: *letter*, **e5**

Alzheimer, The −48 CT polymorphism in the presenilin 1 promoter is associated with an increased risk of developing Alzheimer's disease and an increased Aβ load in brain, **353**

Alzheimer's disease, Genetic association of an *LBP-1c/CP2/LSF* gene polymorphism with late onset Alzheimer's disease, **232**

amniocentesis, Pregnancy outcome and long term prognosis in 868 children born after second trimester amniocentesis for maternal serum positive triple test screening and normal prenatal karyotype: *letter*, **336**

Anderson-Fabry disease, Anderson-Fabry disease: clinical manifestations and impact of disease in a cohort of 60 obligate carrier females: *letter*, **769**

Anderson-Fabry disease: clinical manifestations and impact of disease in a cohort of 98 hemizygous males, **750**

androgen receptor, The androgen receptor and DXS15-134 markers show a high rate of discordance for germline X chromosome inactivation: *letter*, **474**

Angelman syndrome, Angelman syndrome phenotype associated with mutations in *MECP2*, a gene encoding a methyl CpG binding protein, **224**

Distinct phenotypes distinguish the molecular classes of Angelman syndrome, **834**

MECP2 mutation in non-fatal, non-progressive encephalopathy in a male, **171**

Molecular characterisation of four cases of intrachromosomal triplication of chromosome 15q11-q14, **26**

***APC* gene,** A silent mutation in exon 14 of the *APC* gene is associated with exon skipping in a FAP family : *letter*, **863**

A novel 3' mutation in the *APC* gene in a family presenting with a desmoid tumour: *letter*, **861**

The Asp1822Val variant of the *APC* gene is a common polymorphism without clinical implications: *letter*, **e33**

APC, Submicroscopic deletions of the *APC* gene: a frequent cause of familial adenomatous polyposis that may be overlooked by conventional mutation scanning: *letter*, **129**

apolipoprotein B-100, High frequency of the *ApoB*-100 R3500Q mutation in Bulgarian hypercholesterolaemic subjects: *letter*, **536**

apoptosis, Wild type huntingtin reduces the cellular toxicity of mutant huntingtin in mammalian cell models of Huntington's disease, **450**

arachnodactyly, Adducted thumb-club foot syndrome in sibs of a consanguineous Austrian family: *letter*, **265**

arterial hypertrophy, Non-invasive evaluation of arterial involvement in patients affected with Fabry disease, **629**

arylsulphatase A, Multiple metachromatic leucodystrophy alleles in an unaffected subject: a case of dispermic chimaerism: *letter*, **e15**

Asp1822Val variant, The Asp1822Val variant of the *APC* gene is a common polymorphism without clinical implications: *letter*, **e33**

aspartoacylase gene, A partial deletion of the aspartoacylase gene is the cause of Canavan disease in a family from Mexico: *letter*, **e9**

atlantoaxial instability, Anauxetic dysplasia, a spondylometaepiphyseal dysplasia with extreme dwarfism: *letter*, **262**

atopy, Recessively inherited lower incisor hypodontia: *letter*, **551**

attention deficit disorder, Chromosome 2 interstitial deletion (del(2)(q14.1q21)) associated with connective tissue laxity and an attention deficit disorder: *letter*, **493**

attitudes, Participation in preconceptional carrier couple screening: characteristics, attitudes, and knowledge of both partners: *letter*, **695**

autism, Molecular characterisation of four cases of intrachromosomal triplication of chromosome 15q11-q14, **26**

autosomal dominant drusen, Molecular genetic heterogeneity in autosomal dominant drusen, **381**

autosomal dominant inheritance, Acromicric dysplasia: long term outcome and evidence of autosomal dominant inheritance, **745**

autosomal dominant polycystic kidney disease, Autosomal dominant polycystic kidney disease unlinked to the *PKD1* and *PKD2* loci presenting as familial cerebral aneurysm: *letter*, **238**

autosomal recessive, Anauxetic dysplasia, a spondylometaepiphyseal dysplasia with extreme dwarfism: *letter*, **262**

autosomal recessive inheritance, Diaphragmatic hernia, hydrocephalus, and cardiac malformations in four pregnancies of a non-consanguineous couple: *letter*, **269**

autosomal recessive non-syndromic hearing loss, Autosomal recessive non-syndromic hearing loss in the Lebanese population: prevalence of the 30delG mutation and report of two novel mutations in the connexin 26 (*GJB2*) gene: *letter*, **e36**

autozygosity mapping, Kufor-Rakeb syndrome, pallido-pyramidal degeneration with supranuclear upgaze paresis and dementia, maps to 1p36, **680**

B cell immunodeficiency, B cell immunodeficiency, distal limb abnormalities, and urogenital malformations in a three generation family: a novel autosomal dominant syndrome?: *letter*, **488**

Bannayan-Riley-Ruvalcaba syndrome, The spectrum and evolution of phenotypic findings in *PTEN* mutation positive cases of Bannayan-Riley-Ruvalcaba syndrome: *letter*, **52**

Bardet-Biedl syndrome, Does Bardet-Biedl syndrome have a characteristic face?: *letter*, **e14**

behavioural problems, Molecular characterisation of a proximal chromosome 18q deletion: *letter*, **128**

Berry aneurysm, Autosomal dominant polycystic kidney disease unlinked to the *PKD1* and *PKD2* loci presenting as familial cerebral aneurysm: *letter*, **238**

Binder syndrome, Absence of learning difficulties in a hyperactive boy with a terminal Xp deletion encompassing the *MRX49* locus: *letter*, **466**

BPES, Identification of a new *TWIST* mutation (7p21) with variable eyelid manifestations supports locus homogeneity of BPES at 3q22: *letter*, **470**

brachycephaly, Adducted thumb-club foot syndrome in sibs of a consanguineous Austrian family: *letter*, **265**

BRCA1, Interpreting epidemiological research: blinded comparison of methods used to estimate the prevalence of inherited mutations in *BRCA1* , **824**

BRCA1 and *BRCA2* mutation status and cancer family history of Danish women affected with multifocal or bilateral breast cancer at a young age, **361**

Identification of a large rearrangement of the *BRCA1* gene using colour bar code on combed DNA in an American breast/ovarian cancer family previously studied by direct sequencing: *letter*, **388**

BRCA2, *BRCA1* and *BRCA2* mutation status and cancer family history of Danish women affected with multifocal or bilateral breast cancer at a young age, **361**

***BRCA2* gene,** De novo recurrent germline mutation of the *BRCA2* gene in a patient with early onset breast cancer, **102**

breast cancer, A community based study on intentions regarding predictive testing for hereditary breast cancer: *letter*, **540**

De novo recurrent germline mutation of the *BRCA2* gene in a patient with early onset breast cancer, **102**

BRCA1 and *BRCA2* mutation status and cancer family history of Danish women affected with multifocal or bilateral breast cancer at a young age, **361**

Risk perception and cancer worry: an exploratory study of the impact of genetic risk counselling in women with a family history of breast cancer: *letter*, **139**

Science or black magic?: *letter*, **e16**

The importance of good images in using hair to screen for breast cancer: *letter*, **e16**

breast/ovarian cancer, Identification of a large rearrangement of the *BRCA1* gene using colour bar code on combed DNA in an American breast/ovarian cancer family previously studied by direct sequencing: *letter*, **388**

c. 505 T/C germline mutation, *VHL* c.505 T>C mutation confers a high age related penetrance but no increased overall mortality, **508**

C282Y, Variation of iron loading expression in C282Y homozygous haemochromatosis probands and sib pairs: *letter*, **632**

***CACNA1A* gene,** Identification of two novel mutations in the *CACNA1A* gene responsible for episodic ataxia type 2: *letter*, **249**

***CACNA1A*,** Predisposing chromosome for spinocerebellar ataxia type 6 (SCA6) in Japanese: *letter*, **328**

CAG repeat, New problems in testing for Huntington's disease: the issue of intermediate and reduced penetrance alleles: *letter*, **e12**

CAG repeats, Twenty CAG repeats are sufficient to cause the SCA6 phenotype: *letter*, **e38**

Canavan disease, A partial deletion of the aspartoacylase gene is the cause of Canavan disease in a family from Mexico: *letter*, **e9**

cancer, Methylation matters: *review*, **285**

Sensitivity and predictive value of criteria for *p53* germline mutation screening: *letter*, **43**

cancer genetics, Interpreting epidemiological research: blinded comparison of methods used to estimate the prevalence of inherited mutations in *BRCA1* , **824**

Oncology nurse training in cancer genetics: *letter*, **691**

cancer predisposition, Peutz-Jeghers families unlinked to *STK11/LKB1* gene mutations are highly predisposed to primitive biliary adenocarcinoma, **356**

cancer risk, Comprehension of cancer risk one and 12 months after predictive genetic testing for hereditary non-polyposis colorectal cancer: *letter*, **787**

candidate gene, Defining the genetic contribution of type 2 diabetes mellitus: *review*, **569**

candidates, Silver-Russell syndrome: a dissection of the genetic aetiology and candidate chromosomal regions: *review*, **810**

carboxyl-terminal propeptide, Disruption of one intra-chain disulphide bond in the carboxyl-terminal propeptide of the proα1(I) chain of type I procollagen permits slow assembly and secretion of overmodified, but stable procollagen trimers and results in mild osteogenesis imperfecta, **443**

carcinogenesis, Two hits revisited again: *review*, **81**

cardiomyopathy, Mutations in *cis* can confound genotype-phenotype correlations in hypertrophic cardiomyopathy: *letter*, **385**

cardiovascular defects, Mutation analysis of *TBX1* in non-deleted patients with features of DGS/VCFS or isolated cardiovascular defects, **e45**

cataract, A unique form of autosomal dominant cataract explained by gene conversion between β-crystallin B2 and its pseudogene: *letter*, **392**

CATCH22, Deletion 22q11 syndrome: acknowledging a lost eponym as we say farewell to an acronym: *letter*, **271**

cathepsin C, Evidence of a founder effect for four cathepsin C gene mutations in Papillon-Lefèvre syndrome patients, **96**

CDG, A broad spectrum of clinical presentations in congenital disorders of glycosylation I: a series of 26 cases, **14**

CENP-C, Alternative centromeric inactivation in a pseudodicentric t(Y;13)(q12;p11. 2) translocation chromosome associated with extreme oligozoospermia: *letter*, **802**

centromere, Alternative centromeric inactivation in a pseudodicentric t(Y;13)(q12;p11.2) translocation chromosome associated with extreme oligozoospermia: *letter*, **802**

***CFTR* gene,** Analysis of the entire coding region of the cystic fibrosis transmembrane regulator gene in neonatal hypertrypsinaemia with normal sweat test: *letter*, **202**

Mutation and haplotype analysis of the *CFTR* gene in atypically mild cystic fibrosis patients from Northern Ireland: *letter*, **136**

CGG repeats, Microdeletion in the *FMR-1* gene: an apparent null allele using routine clinical PCR amplification: *letter*, **624**

characteristic face, Does Bardet-Biedl syndrome have a characteristic face?: *letter*, **e14**

checklist, Clinical studies on submicroscopic subtelomeric rearrangements: a checklist, **145**

children, Predictive genetic testing in children and adults: a study of emotional impact, **519**

chimaerism, Multiple metachromatic leucodystrophy alleles in an unaffected subject: a case of dispermic chimaerism: *letter*, **e15**

Chinese, A Chinese adult onset type II citrullinaemia patient with 851del4/1638ins23 mutations in the *SLC25A13* gene: *letter*, **e23**

Fibrinogen genotypes (α and β) are associated with plasma fibrinogen levels in Chinese: *letter*, **e31**

chondrosarcoma, Genotype-phenotype correlation in hereditary multiple exostoses, **430**

chromosome 1, Genetic heterogeneity and exclusion of a modifying locus at 17p11.2-p11.1 in Finnish families with van der Woude syndrome: *letter*, **198**

High frequencies of ICF syndrome-like pericentromeric heterochromatin decondensation and breakage in chromosome 1 in a chorionic villus sample : *letter*, **882**

Supernumerary marker chromosome (1) of paternal origin and maternal uniparental disomy 1 in a developmentally delayed child: *letter*, **885**

chromosome 11, A syndrome of overgrowth and acromegaloidism with normal growth hormone secretion is associated with chromosome 11 pericentric inversion: *letter*, **338**

Interstitial deletion of chromosome 11 (q22.3-q23.2) in a boy with mild developmental delay: *letter*, **621**

chromosome 11q14, Evidence of a founder effect for four cathepsin C gene mutations in Papillon-Lefèvre syndrome patients, **96**

chromosome 12, Satellites on the terminal short arm of chromosome 12 (12ps), inherited through several generations in three families: a new variant without phenotypic effect: *letter*, **723**

chromosome 14q, Hereditary spastic paraplegia linked to chromosome 14q11-q21: reduction of the SPG3 locus interval from 5.3 to 2.7 cM: *letter*, **e39**

chromosome 15 triplication, Molecular characterisation of four cases of intrachromosomal triplication of chromosome 15q11-q14, **26**

chromosome 15q26, Localisation of a gene for an autosomal recessive syndrome of macrocephaly, multiple epiphyseal dysplasia, and distinctive facies to chromosome 15q26, **369**

chromosome 17, Genetic heterogeneity and exclusion of a modifying locus at 17p11.2-p11.1 in Finnish families with van der Woude syndrome: *letter*, **198**

chromosome 19, Mosaic terminal del(19)(q13.33:) in a girl with seizures and mental retardation: *letter*, **e2**

chromosome 1p36, Kufor-Rakeb syndrome, pallido-pyramidal degeneration with supranuclear upgaze paresis and dementia, maps to 1p36, **680**

chromosome 2 deletion, A transmitted deletion of 2q13 to 2q14.1 causes no phenotypic abnormalities: *letter*, **125**

chromosome 2 interstitial deletion, Chromosome 2 interstitial deletion (del(2)(q14.1q21)) associated with connective tissue laxity and an attention deficit disorder: *letter*, **493**

chromosome 2p16, Segmental uniparental isodisomy (UPD) for 2p16 without clinical symptoms: implications for UPD and other genetic studies of chromosome 2: *letter*, **106**

chromosome 3, A dominant gene for developmental dyslexia on chromosome 3, **658**

chromosome 4, An epidemiological study of Wolf-Hirschhorn syndrome: life expectancy and cause of mortality, **674**

Ellis-van Creveld syndrome resulting from segmental uniparental disomy of chromosome 4: *letter*, **e18**

chromosome 6p, Suggestive linkage of situs inversus and other left-right axis anomalies to chromosome 6p: *letter*, **182**

chromosome 7, Split hand/split foot malformation associated with sensorineural deafness, inner and middle ear malformation, hypodontia, congenital vertical talus, and deletion of eight microsatellite markers in 7q21.1-q21.3: *letter*, **405**

chromosome 7q36, A novel acropectoral syndrome maps to chromosome 7q36, **304**

chromosome aberrations, High resolution comparative genomic hybridisation in clinical cytogenetics, **740**

chromosome analysis, High resolution comparative genomic hybridisation in clinical cytogenetics, **740**

chromosome deletion, Clinical studies on submicroscopic subtelomeric rearrangements: a checklist, **145**

chromosome microdissection, Supernumerary marker chromosome (1) of paternal origin and maternal uniparental disomy 1 in a developmentally delayed child : *letter*, **885**

chromosome Xptel-Xp22.3, Absence of learning difficulties in a hyperactive boy with a terminal Xp deletion encompassing the *MRX49* locus: *letter*, **466**

chronic idiopathic pancreatitis, Low prevalence of *SPINK1* gene mutations in adult patients with chronic idiopathic pancreatitis: *letter*, **243**

chronic progressive external ophthalmoplegia, Functional characterisation of mitochondrial $tRNA^{Tyr}$ mutation (5877G→A) associated with familial chronic progressive external ophthalmoplegia: *letter*, **703**

circadian rhythms, β_1-adrenergic antagonists improve sleep and behavioural disturbances in a circadian disorder, Smith-Magenis syndrome, **586**

cirrhosis, The mannose binding lectin gene influences the severity of chronic liver disease in cystic fibrosis, **310**
citrullinaemia, A Chinese adult onset type II citrullinaemia patient with 851del4/1638ins23 mutations in the *SLC25A13* gene: *letter,* **e23**
cleft lip and palate, Genetic heterogeneity and exclusion of a modifying locus at 17p11.2-p11.1 in Finnish families with van der Woude syndrome: *letter,* **198**
clinical features, *SHOX* haploinsufficiency and overdosage: impact of gonadal function status: *review,* **1**
clinical heterogeneity, Clinical heterogeneity in lymphoedema-distichiasis with *FOXC2* truncating mutations, **761**
clinical preselection, Clinical studies on submicroscopic subtelomeric rearrangements: a checklist, **145**
clinical-functional correlation, Ectodermal dysplasias: a new clinical-genetic classification: *review,* **579**
club foot, Adducted thumb-club foot syndrome in sibs of a consanguineous Austrian family: *letter,* **265**
Sulphate transporter gene mutations in apparently isolated club foot: *letter,* **191**
c-MYC, Identification of a transcriptionally compromised allele of *c-MYC* in a North American family: *letter,* **47**
coeliac disease, Coeliac disease in Williams syndrome, **767**
collagen mutations, Homozygosity for a splice site mutation of the *COL1A2* gene yields a non-functional proα2(I) chain and an EDS/OI clinical phenotype: *letter,* **132**
colorectal cancer, Germline mutation analysis of the transforming growth factor β receptor type II (*TGFBR2*) and E-cadherin (*CDH1*) genes in early onset and familial colorectal cancer: *letter,* **e7**
Hereditary and somatic DNA mismatch repair gene mutations in sporadic endometrial carcinoma: *letter,* **461**
Screening families with endometrial and colorectal cancers for germline mutations: *letter,* **e29**
colour bar code, Identification of a large rearrangement of the *BRCA1* gene using colour bar code on combed DNA in an American breast/ovarian cancer family previously studied by direct sequencing: *letter,* **388**
community genetics, A community based study on intentions regarding predictive testing for hereditary breast cancer: *letter,* **540**
comparative genomic hybridisation, De novo terminal deletion of chromosome 15q26.1 characterised by comparative genomic hybridisation and FISH with locus specific probes: *letter,* **617**
High resolution comparative genomic hybridisation in clinical cytogenetics, **740**
complex chromosomal rearrangement (CCR), Molecular cytogenetic characterisation of a complex 46,XY,t(7;8;11;13) chromosome rearrangement in a patient with Moebius syndrome: *letter,* **117**
cone-rod dystrophy, Clustering and frequency of mutations in the retinal guanylate cyclase (*GUCY2D*) gene in patients with dominant cone-rod dystrophies: *letter,* **611**
congenital heart disease, A region of homozygosity within 22q11.2 associated with congenital heart disease: recessive DiGeorge/velocardiofacial syndrome?: *letter,* **533**
congenital muscular dystrophy, The expanding phenotype of laminin α2 chain (merosin) abnormalities: case series and review: *review,* **649**
connective tissue laxity, Chromosome 2 interstitial deletion (del(2)(q14.1q21)) associated with connective tissue laxity and an attention deficit disorder: *letter,* **493**
connexin 26, A common founder for the 35delG *GJB2* gene mutation in connexin 26 hearing impairment, **515**
Recurrent mutations in the deafness gene *GJB2* (connexin 26) in British Asian families: *letter,* **530**
connexin 26 gene, Autosomal recessive non-syndromic hearing loss in the Lebanese population: prevalence of the 30delG mutation and report of two novel mutations in the connexin 26 (*GJB2*) gene: *letter,* **36**
connexin-26, Genetic analysis of the connexin-26 M34T variant: identification of genotype M34T/M34T segregating with mild-moderate non-syndromic sensorineural hearing loss, **20**
Genetic analysis of the connexin-26 M34T variant: *letter,* **e24**
contraction, Trinucleotide repeat contraction: a pitfall in prenatal diagnosis of myotonic dystrophy : *letter,* **850**
corneal dermoids, Potential mapping of corneal dermoids to Xq24-qter: *letter,* **719**
corneal nerves, Familial medullary thyroid carcinoma and prominent corneal nerves associated with the germline V804M and V778I mutations on the same allele of *RET*: *letter,* **784**
Cowden syndrome, Male breast cancer in Cowden syndrome patients with germline *PTEN* mutations, **159**
The spectrum and evolution of phenotypic findings in *PTEN* mutation positive cases of Bannayan-Riley-Ruvalcaba syndrome: *letter,* **52**
COX deficiency, Mutations in *SURF1* are not specifically associated with Leigh syndrome: *letter,* **109**
cri du chat syndrome, Clinical and molecular characterisation of 80 patients with 5p deletion: genotype-phenotype correlation, **151**
cryptic telomeric rearrangements, Cryptic telomeric rearrangements in subjects with mental retardation associated with dysmorphism and congenital malformations: *letter,* **417**
crystallin, A unique form of autosomal dominant cataract explained by gene conversion between β-crystallin B2 and its pseudogene: *letter,* **392**
***CTNS* mutation,** A G339R mutation in the *CTNS* gene is a common cause of nephropathic cystinosis in the south western Ontario Amish Mennonite population: *letter,* **615**
cultural competence, Cultural aspects of cancer genetics: setting a research agenda: *review,* **425**
culture, Cultural aspects of cancer genetics: setting a research agenda: *review,* **425**
CYP1B1, Phenotypic heterogeneity of *CYP1B1*: mutations in a patient with Peters' anomaly: *letter,* **324**
cysteine substitution, A cysteine substitution in the zona pellucida domain of α-tectorin results in autosomal dominant, postlingual, progressive, mid frequency hearing loss in a Spanish family: *letter,* **e13**
cystic fibrosis, A novel 3600+11.5 kb C>G homozygous splicing mutation in a black African, consanguineous CF family: *letter,* **e4**
Analysis of the entire coding region of the cystic fibrosis transmembrane regulator gene in neonatal hypertrypsinaemia with normal sweat test: *letter,* **202**
Cystic fibrosis patients with the 3272-26A>G splicing mutation have milder disease than F508del homozygotes: a large European study: *letter,* **777**
Mutation and haplotype analysis of the *CFTR* gene in atypically mild cystic fibrosis patients from Northern Ireland: *letter,* **136**
Participation in preconceptional carrier couple screening: characteristics, attitudes, and knowledge of both partners: *letter,* **695**
The mannose binding lectin gene influences the severity of chronic liver disease in cystic fibrosis, **310**
cystinosin, A G339R mutation in the *CTNS* gene is a common cause of nephropathic cystinosis in the south western Ontario Amish Mennonite population: *letter,* **615**

Dandy-Walker malformation, Cytogenetic and molecular study of a jumping translocation in a baby with Dandy-Walker malformation: *letter,* **67**
de novo mutation, De novo recurrent germline mutation of the *BRCA2* gene in a patient with early onset breast cancer, **102**
deafness, Nijmegen breakage syndrome in a Dutch patient not resulting from a defect in *NBS1: letter,* **e19**
Recurrent mutations in the deafness gene *GJB2* (connexin 26) in British Asian families: *letter,* **530**
DEFECT 11, DEFECT 11 syndrome associated with agenesis of the corpus callosum: *letter,* **e5**
deletion, A frameshift mitochondrial complex I gene mutation in a patient with dystonia and cataracts: is the mutation pathogenic?: *letter,* **58**
A simple non-isotopic method to show pitfalls during mutation analysis of the glucocerebrosidase gene: *letter,* **e34**
deletion 22q11 syndrome, Deletion 22q11 syndrome: acknowledging a lost eponym as we say farewell to an acronym: *letter,* **271**
dementia, Genetic association of an *LBP-1c/CP2/LSF* gene polymorphism with late onset Alzheimer's disease, **232**
desmoid tumour, A novel 3' mutation in the *APC* gene in a family presenting with a desmoid tumour: *letter,* **861**
developmental delay, Interstitial deletion of 3p22.2-p24.2: the first reported case: *letter,* **349**
Interstitial deletion of chromosome 11 (q22.3-q23.2) in a boy with mild developmental delay: *letter,* **621**
DFNA9, A common ancestor for *COCH* related cochleovestibular (DFNA9) patients in Belgium and The Netherlands bearing the P51S mutation: *letter,* **61**
DFNB10, Novel mutations of *TMPRSS3* in four DFNB8/B10 families segregating congenital autosomal recessive deafness: *letter,* **396**
DFNB8, Novel mutations of *TMPRSS3* in four DFNB8/B10 families segregating congenital autosomal recessive deafness: *letter,* **396**
DHPLC, Recurrent mutations in the deafness gene *GJB2* (connexin 26) in British Asian families: *letter,* **530**
diabetes mellitus, Wolfram syndrome: a clinical and molecular genetic analysis: *letter,* **e37**
diabetes mellitus type 2, Defining the genetic contribution of type 2 diabetes mellitus: *review,* **569**
diaphragmatic hernia, Congenital diaphragmatic hernia and interstitial deletion of chromosome 3: *letter,* **556**
Diaphragmatic hernia, hydrocephalus, and cardiac malformations in four pregnancies of a non-consanguineous couple: *letter,* **269**
DiGeorge syndrome, Mutation analysis of *TBX1* in non-deleted patients with features of DGS/VCFS or isolated cardiovascular defects, **e45**
DiGeorge/velocardiofacial syndrome, A region of homozygosity within 22q11.2 associated with congenital heart disease: recessive DiGeorge/velocardiofacial syndrome?: *letter,* **533**
disease manifestations, Anderson-Fabry disease: clinical manifestations and impact of disease in a cohort of 60 obligate carrier females: *letter,* **769**
distal arthrogryposis, Distal arthrogryposis: a new type with distinct facial appearance and absent teeth: *letter,* **e22**

distal limb abnormalities, B cell immunodeficiency, distal limb abnormalities, and urogenital malformations in a three generation family: a novel autosomal dominant syndrome?: *letter,* **488**
distichiasis, Clinical heterogeneity in lymphoedema-distichiasis with *FOXC2* truncating mutations, **761**
distinctive facies, Localisation of a gene for an autosomal recessive syndrome of macrocephaly, multiple epiphyseal dysplasia, and distinctive facies to chromosome 15q26, **369**
DNA methylation, Distinct phenotypes distinguish the molecular classes of Angelman syndrome, **834**
DNA mismatch repair, Hereditary and somatic DNA mismatch repair gene mutations in sporadic endometrial carcinoma: *letter,* **461**
dolichyl-phosphate glucose:mannose 9 N-acetylglycosamine 2 glucosyltransferase, A broad spectrum of clinical presentations in congenital disorders of glycosylation I: a series of 26 cases, **14**
dosage quotients, Submicroscopic deletions of the *APC* gene: a frequent cause of familial adenomatous polyposis that may be overlooked by conventional mutation scanning: *letter,* **129**
Down syndrome, A novel mutation in the endothelin B receptor gene in a patient with Shah-Waardenburg syndrome and Down syndrome: *letter,* **646**
Clinical and cytogenetic characterisation of a patient with Down syndrome resulting from a 21q22.1→qter duplication: *letter,* **73**
DTDST, Sulphate transporter gene mutations in apparently isolated club foot: *letter,* **191**
duplication, A comparison of methods for gene dosage analysis in HMSN type 1, **90**
duplication 7p, Pure partial 7p trisomy including the *TWIST*, *HOXA*, and *GLI3* genes: *letter,* **178**
DXS15-134, The androgen receptor and DXS15-134 markers show a high rate of discordance for germline X chromosome inactivation: *letter,* **474**
dyschondrosteosis, *SHOX* point mutations in dyschondrosteosis: *letter,* **323**
dysmorphism, High resolution comparative genomic hybridisation in clinical cytogenetics, **740**
***DYT1* gene,** No evidence of allelic heterogeneity in the *DYT1* gene of European patients with early onset torsion dystonia: *letter,* **e35**

E-cadherin, Germline mutation analysis of the transforming growth factor β receptor type II (*TGFBR2*) and E-cadherin (*CDH1*) genes in early onset and familial colorectal cancer: *letter,* **e7**
E-cadherin, *E-cadherin* is not frequently mutated in hereditary gastric cancer: *letter,* **49**
ectodermal dysplasia, Ectodermal dysplasias: a new clinical-genetic classification: *review,* **579**
ectodermal structural proteins, Ectodermal dysplasias: a new clinical-genetic classification: *review,* **579**
Ehlers-Danlos syndrome, Homozygosity for a splice site mutation of the *COL1A2* gene yields a non-functional proα2(I) chain and an EDS/OI clinical phenotype: *letter,* **132**
electronic publication, *JMG* Online: exploiting the potential of electronic publication and manuscript submission: *editorial,* **809**
Ellis-van Creveld syndrome, Ellis-van Creveld syndrome resulting from segmental uniparental disomy of chromosome 4: *letter,* **e18**
***ELOVL4*,** Evaluation of the *ELOVL4* gene in families with retinitis pigmentosa linked to the *RP25* locus: *letter,* **478**
emotional impact, Predictive genetic testing in children and adults: a study of emotional impact, **519**
encephalocele, Laryngeal atresia, encephalocele, and limb deformities (LEL): a possible new syndrome: *letter,* **420**
encephalomyopathy, Mitochondrial DNA inheritance in patients with deleted mtDNA: *letter,* **e28**
encephalopathy, *MECP2* mutation in non-fatal, non-progressive encephalopathy in a male, **171**
endometrial cancer, Hereditary and somatic DNA mismatch repair gene mutations in sporadic endometrial carcinoma: *letter,* **461**
Screening families with endometrial and colorectal cancers for germline mutations: *letter,* **e29**
endothelin 3 mutation, A heterozygous endothelin 3 mutation in Waardenburg-Hirschsprung disease: is there a dosage effect of *EDN3*/*EDNRB* gene mutations on neurocristopathy phenotypes?: *letter,* **205**
endothelin B receptor gene, A novel mutation in the endothelin B receptor gene in a patient with Shah-Waardenburg syndrome and Down syndrome: *letter,* **646**
enteropathy, Novel mutations of *FOXP3* in two Japanese patients with immune dysregulation, polyendocrinopathy, enteropathy, X linked syndrome (IPEX): *letter,* **874**
episodic ataxia type 2, Identification of two novel mutations in the *CACNA1A* gene responsible for episodic ataxia type 2: *letter,* **249**
epistasis, Identification of a novel psoriasis susceptibility locus at 1p and evidence of epistasis between *PSORS1* and candidate loci, **7**
epithelial-mesenchymal interaction, Ectodermal dysplasias: a new clinical-genetic classification: *review,* **579**
epoxide hydrolase, A polymorphism in the gene for microsomal epoxide hydrolase is associated with pre-eclampsia, **234**
ethics, The whole truth and nothing but the truth, but what is the truth?, **39**
exomphalos, A new case of exomphalos, short limbs, and macrogonadism syndrome: *letter,* **e8**
exon skipping, A silent mutation in exon 14 of the *APC* gene is associated with exon skipping in a FAP family : *letter,* **863**
***EXT1*,** Genotype-phenotype correlation in hereditary multiple exostoses, **430**
***EXT2*,** Genotype-phenotype correlation in hereditary multiple exostoses, **430**
eyelids, Identification of a new *TWIST* mutation (7p21) with variable eyelid manifestations supports locus homogeneity of BPES at 3q22: *letter,* **470**

F syndrome, A novel acropectoral syndrome maps to chromosome 7q36, **304**
Fabry disease, Non-invasive evaluation of arterial involvement in patients affected with Fabry disease, **629**
familial adenomatous polyposis, A silent mutation in exon 14 of the *APC* gene is associated with exon skipping in a FAP family: *letter,* **863**
Predictive genetic testing in children and adults: a study of emotional impact, **519**
Submicroscopic deletions of the *APC* gene: a frequent cause of familial adenomatous polyposis that may be overlooked by conventional mutation scanning: *letter,* **129**
The Asp1822Val variant of the *APC* gene is a common polymorphism without clinical implications: *letter,* **e33**
familial hypertrophic cardiomyopathy, Myosin light chain mutations in familial hypertrophic cardiomyopathy: phenotypic presentation and frequency in Danish and South African populations: *letter,* **e43**
familial hypertrophic myopathy, Development and application of linkage analysis in genetic diagnosis of familial hypertrophic cardiomyopathy: *letter,* **193**
familial non-medullary thyroid cancer, Familial non-medullary thyroid cancer in Iceland: *letter,* **189**
familial renal cell carcinoma, Association of a novel constitutional translocation t(1q;3q) with familial renal cell carcinoma, **165**
family history, Cultural aspects of cancer genetics: setting a research agenda: *review,* **425**
Risk perception and cancer worry: an exploratory study of the impact of genetic risk counselling in women with a family history of breast cancer: *letter,* **139**
fibrinogen genotypes, Fibrinogen genotypes (α and β) are associated with plasma fibrinogen levels in Chinese: *letter,* **31**
FISH, Clinical and cytogenetic characterisation of a patient with Down syndrome resulting from a 21q22.1→qter duplication: *letter,* **73**
Clinical and molecular characterisation of 80 patients with 5p deletion: genotype-phenotype correlation, **151**
fluorescence in situ hybridisation, A syndrome of overgrowth and acromegaloidism with normal growth hormone secretion is associated with chromosome 11 pericentric inversion: *letter,* **338**
fluorescence in situ hybridisation (FISH), Molecular cytogenetic characterisation of a complex 46,XY,t(7;8;11;13) chromosome rearrangement in a patient with Moebius syndrome: *letter,* **117**
***FMR-1* gene,** Microdeletion in the *FMR-1* gene: an apparent null allele using routine clinical PCR amplification: *letter,* **624**
forkhead gene, Clinical heterogeneity in lymphoedema-distichiasis with *FOXC2* truncating mutations, **761**
founder effect, A common founder for the 35delG *GJB2* gene mutation in connexin 26 hearing impairment, **515**
Evidence of a founder effect for four cathepsin C gene mutations in Papillon-Lefèvre syndrome patients, **96**
Founder effect in multiple endocrine neoplasia type 1 (MEN 1) in Finland: *letter,* **185**
founder mutations, Dutch patients with glycogen storage disease type II show common ancestry for the 525delT and del exon 18 mutations: *letter,* **527**
FOXP3 gene, Novel mutations of *FOXP3* in two Japanese patients with immune dysregulation, polyendocrinopathy, enteropathy, X linked syndrome (IPEX): *letter,* **874**
fragile X syndrome, Microdeletion in the *FMR-1* gene: an apparent null allele using routine clinical PCR amplification: *letter,* **624**
Friderici, A case of Roberts syndrome described in 1737: *letter,* **565**
Fryns syndrome, Diaphragmatic hernia, hydrocephalus, and cardiac malformations in four pregnancies of a non-consanguineous couple: *letter,* **269**

GATA3, GATA3 abnormalities and the phenotypic spectrum of HDR syndrome, **374**
Gaucher disease, A simple non-isotopic method to show pitfalls during mutation analysis of the glucocerebrosidase gene: *letter,* **e34**
***GDNF*,** *GDNF* as a candidate modifier in a type 1 neurofibromatosis (NF1) enteric phenotype, **638**
geleophysic dysplasia, Acromicric dysplasia: long term outcome and evidence of autosomal dominant inheritance, **745**
gene conversion, A unique form of autosomal dominant cataract explained by gene conversion between β-crystallin B2 and its pseudogene: *letter,* **392**
Limited contribution of interchromosomal gene conversion to *NF1* gene mutation: *letter,* **481**

gene dosage, A comparison of methods for gene dosage analysis in HMSN type 1, **90**

gene mutation, A missense mutation in the *SEDL* gene results in delayed onset of X linked spondyloepiphyseal dysplasia in a large pedigree: *letter,* **409**

gene silencing, A majority of fragile X males with methylated, full mutation alleles have significant levels of *FMR1* messenger RNA, **453**

genes, Mutations in *cis* can confound genotype-phenotype correlations in hypertrophic cardiomyopathy: *letter,* **385**

genetic counselling, A community based study on intentions regarding predictive testing for hereditary breast cancer: *letter,* **540**

A simple non-isotopic method to show pitfalls during mutation analysis of the glucocerebrosidase gene: *letter,* **e34**

Presymptomatic testing in myotonic dystrophy: genetic counselling approaches : *letter,* **846**

Risk perception and cancer worry: an exploratory study of the impact of genetic risk counselling in women with a family history of breast cancer: *letter,* **139**

genetic factors, Defining the genetic contribution of type 2 diabetes mellitus: *review,* **569**

genetic heterogeneity, Peutz-Jeghers families unlinked to *STK11/LKB1* gene mutations are highly predisposed to primitive biliary adenocarcinoma, **356**

genetic polymorphism, A polymorphism in the gene for microsomal epoxide hydrolase is associated with pre-eclampsia, **234**

genetic testing, Hair roots as the ideal source of mRNA for genetic testing: *letter,* **e20**

Predictive genetic testing in children and adults: a study of emotional impact, **519**

genetics, Hirschsprung disease, associated syndromes, and genetics: a review: *review,* **729**

The whole truth and nothing but the truth, but what is the truth?, **39**

genodermatoses, Keratosis pilaris/ulerythema ophryogenes and 18p deletion: is it possible that the *LAMA1* gene is involved?: *letter,* **127**

genome screen, Defining the genetic contribution of type 2 diabetes mellitus: *review,* **569**

genomic imprinting, Complex and segmental uniparental disomy (UPD): review and lessons from rare chromosomal complements: *review,* **497**

genotype-phenotype, Mutations in *cis* can confound genotype-phenotype correlations in hypertrophic cardiomyopathy: *letter,* **385**

genotype-phenotype correlation, Cystic fibrosis patients with the 3272-26A>G splicing mutation have milder disease than F508del homozygotes: a large European study: *letter,* **777**

genotype-phenotype correlations, Distinct phenotypes distinguish the molecular classes of Angelman syndrome, **834**

Gilbert's syndrome, Interaction of coding region mutations and the Gilbert-type promoter abnormality of the *UGT1A1* gene causes moderate degrees of unconjugated hyperbilirubinaemia and may lead to neonatal kernicterus: *letter,* **244**

GJB2, A common founder for the 35delG *GJB2* gene mutation in connexin 26 hearing impairment, **515**

Autosomal recessive non-syndromic hearing loss in the Lebanese population: prevalence of the 30delG mutation and report of two novel mutations in the connexin 26 (*GJB2*) gene: *letter,* **e36**

Genetic analysis of the connexin-26 M34T variant: identification of genotype M34T/M34T segregating with mild-moderate non-syndromic sensorineural hearing loss, **20**

Genetic analysis of the connexin-26 M34T variant: *letter,* **e24**

glaucoma, Ocular malformations, postaxial polydactyly, and delayed intramembranous ossification: a new autosomal dominant condition: *letter,* **547**

Phenotypic heterogeneity of *CYP1B1*: mutations in a patient with Peters' anomaly: *letter,* **324**

glioblastoma, A distinct splice form of *APC* is highly expressed in neurones but not commonly mutated in neuroepithelial tumours: *letter,* **257**

GLRA1, Mutations in the glycine receptor α1 subunit (*GLRA1*) gene in hereditary hyperekplexia pedigrees: evidence for non-penetrance of mutation Y279C: *letter,* **e17**

glucocerebrosidase, A simple non-isotopic method to show pitfalls during mutation analysis of the glucocerebrosidase gene: *letter,* **e34**

glycine α1 subunit, Mutations in the glycine receptor α1 subunit (*GLRA1*) gene in hereditary hyperekplexia pedigrees: evidence for non-penetrance of mutation Y279C: *letter,* **e17**

glycogenosis, Dutch patients with glycogen storage disease type II show common ancestry for the 525delT and del exon 18 mutations: *letter,* **527**

glycoprotein IIIa, Thrombophilic polymorphisms in pre-eclampsia: altered frequency of the functional 98C>T polymorphism of glycoprotein IIIa: *letter,* **775**

growth retardation, A case of ring chromosome 2 with growth retardation, mild dysmorphism, and microdeletion of 2p detected using FISH: *letter,* **e32**

Identification of interstitial maternal uniparental disomy (UPD) (14) and complete maternal UPD(20) in a cohort of growth retarded patients, **86**

GTPase activating protein homology region, A novel missense mutation in the GTPase activating protein homology region of *TSC2* in two large families with tuberous sclerosis complex: *letter,* **347**

guanylate cyclase, Clustering and frequency of mutations in the retinal guanylate cyclase (*GUCY2D*) gene in patients with dominant cone-rod dystrophies: *letter,* **611**

GUCY2D, Clustering and frequency of mutations in the retinal guanylate cyclase (*GUCY2D*) gene in patients with dominant cone-rod dystrophies: *letter,* **611**

haemochromatosis, Classification and genetic features of neonatal haemochromatosis: a study of 27 affected pedigrees and molecular analysis of genes implicated in iron metabolism, **599**

Variation of iron loading expression in C282Y homozygous haemochromatosis probands and sib pairs: *letter,* **632**

haemophagocytic lymphohistiocytosis, Six novel mutations in the *PRF1* gene in children with haemophagocytic lymphohistiocytosis: *letter,* **643**

hair, Science or black magic?: *letter,* **e16**

The importance of good images in using hair to screen for breast cancer: *letter,* **e16**

hair roots, Hair roots as the ideal source of mRNA for genetic testing: *letter,* **e20**

haplotype, Dutch patients with glycogen storage disease type II show common ancestry for the 525delT and del exon 18 mutations: *letter,* **527**

haptoglobin genotype, Haptoglobin genotype as a risk factor for postmenopausal osteoporosis: *letter,* **636**

HDR syndrome, GATA3 abnormalities and the phenotypic spectrum of HDR syndrome, **374**

hearing impairment, A common ancestor for *COCH* related cochleovestibular (DFNA9) patients in Belgium and The Netherlands bearing the P51S mutation: *letter,* **61**

Prevalence of mitochondrial DNA mutations in childhood/congenital onset non-syndromal sensorineural hearing impairment, **229**

hearing loss, Genetic analysis of the connexin-26 M34T variant: identification of genotype M34T/M34T segregating with mild-moderate non-syndromic sensorineural hearing loss, **20**

Genetic analysis of the connexin-26 M34T variant: *letter,* **e24**

HELLP syndrome, A polymorphism in the gene for microsomal epoxide hydrolase is associated with pre-eclampsia, **234**

hereditary cancer, Cultural aspects of cancer genetics: setting a research agenda: *review,* **425**

hereditary gastric cancer, *E-cadherin* is not frequently mutated in hereditary gastric cancer: *letter,* **49**

hereditary hyperekplexia, Mutations in the glycine receptor α1 subunit (*GLRA1*) gene in hereditary hyperekplexia pedigrees: evidence for non-penetrance of mutation Y279C: *letter,* **e17**

hereditary multiple exostoses, Genotype-phenotype correlation in hereditary multiple exostoses, **430**

hereditary non-polyposis colorectal cancer, Atypical HNPCC owing to *MSH6* germline mutations: analysis of a large Dutch pedigree, **318**

Comprehension of cancer risk one and 12 months after predictive genetic testing for hereditary non-polyposis colorectal cancer: *letter,* **787**

hereditary spastic paraplegia, Hereditary spastic paraplegia linked to chromosome 14q11-q21: reduction of the SPG3 locus interval from 5.3 to 2.7 cM: *letter,* **e39**

heteroplasmy, Heterogeneous tissue distribution of a mitochondrial DNA polymorphism in heteroplasmic subjects without mitochondrial disorders, **312**

heterotaxy, Suggestive linkage of situs inversus and other left-right axis anomalies to chromosome 6p: *letter,* **182**

Hirschsprung disease, Hirschsprung disease, associated syndromes, and genetics: a review: *review,* **729**

history, A case of Roberts syndrome described in 1737: *letter,* **565**

HMSN, A comparison of methods for gene dosage analysis in HMSN type 1, **90**

HNPCC, Screening families with endometrial and colorectal cancers for germline mutations: *letter,* **e29**

Holt-Oram syndrome, Detection of a large *TBX5* deletion in a family with Holt-Oram syndrome, **e44**

homoplasmic 3316G→A mutation, Homoplasmic 3316G→A in the *ND1* gene of the mitochondrial genome: a pathogenic mutation or a neutral polymorphism?: *letter,* **e10**

homozygosity, Molecular and clinical study of two myotonic dystrophy homozygotes: *letter,* **e40**

human chromosome 21, Clinical and cytogenetic characterisation of a patient with Down syndrome resulting from a 21q22.1→qter duplication: *letter,* **73**

human papillomavirus, Role of *TP53* P72R polymorphism in human papillomavirus associated premalignant laryngeal neoplasm: *letter,* **327**

huntingtin, Wild type huntingtin reduces the cellular toxicity of mutant huntingtin in mammalian cell models of Huntington's disease, **450**

Huntington's disease, New problems in testing for Huntington's disease: the issue of intermediate and reduced penetrance alleles: *letter*, **e12**
Psychological studies in Huntington's disease: making up the balance: *letter*, **852**
The whole truth and nothing but the truth, but what is the truth?, **39**
Wild type huntingtin reduces the cellular toxicity of mutant huntingtin in mammalian cell models of Huntington's disease, **450**
HVR2, Heterogeneous tissue distribution of a mitochondrial DNA polymorphism in heteroplasmic subjects without mitochondrial disorders, **312**
hydrocephalus, A novel germline mutation of the *PTEN* gene in a patient with macrocephaly, ventricular dilatation, and features of VATER association, **820**
Diaphragmatic hernia, hydrocephalus, and cardiac malformations in four pregnancies of a non-consanguineous couple: *letter*, **269**
hyperactivity, Absence of learning difficulties in a hyperactive boy with a terminal Xp deletion encompassing the *MRX49* locus: *letter*, **466**
hyperbilirubinaemia, Interaction of coding region mutations and the Gilbert-type promoter abnormality of the *UGT1A1* gene causes moderate degrees of unconjugated hyperbilirubinaemia and may lead to neonatal kernicterus: *letter*, **244**
hypercholesterolaemia, High frequency of the *ApoB*-100 R3500Q mutation in Bulgarian hypercholesterolaemic subjects: *letter*, **536**
hypodontia, Recessively inherited lower incisor hypodontia: *letter*, **551**

Iceland, Familial non-medullary thyroid cancer in Iceland: *letter*, **189**
ICF syndrome, High frequencies of ICF syndrome-like pericentromeric heterochromatin decondensation and breakage in chromosome 1 in a chorionic villus sample : *letter*, **882**
IGF1R, De novo terminal deletion of chromosome 15q26.1 characterised by comparative genomic hybridisation and FISH with locus specific probes: *letter*, **617**
imprinting, Maternally inherited duplication of the possible imprinted 14q31 region: *letter*, **343**
Silver-Russell syndrome: a dissection of the genetic aetiology and candidate chromosomal regions: *review*, **810**
in situ hybridisation, De novo terminal deletion of chromosome 15q26.1 characterised by comparative genomic hybridisation and FISH with locus specific probes: *letter*, **617**
incidence, Population prevalence and estimated birth incidence and mortality rate for people with Prader-Willi syndrome in one UK Health Region: *letter*, **792**
informed consent, The whole truth and nothing but the truth, but what is the truth?, **39**
insertion, A frameshift mitochondrial complex I gene mutation in a patient with dystonia and cataracts: is the mutation pathogenic?: *letter*, **58**
interstitial deletion, Interstitial deletion of chromosome 11 (q22.3-q23.2) in a boy with mild developmental delay: *letter*, **621**
interstitial deletion of chromosome 3, Congenital diaphragmatic hernia and interstitial deletion of chromosome 3: *letter*, **556**
intrinsic intestinal dysmotility, *GDNF* as a candidate modifier in a type 1 neurofibromatosis (NF1) enteric phenotype, **638**
inversion, A syndrome of overgrowth and acromegaloidism with normal growth hormone secretion is associated with chromosome 11 pericentric inversion: *letter*, **338**
IPEX syndrome, Novel mutations of *FOXP3* in two Japanese patients with immune dysregulation, polyendocrinopathy, enteropathy, X linked syndrome (IPEX): *letter*, **874**
iron, Haptoglobin genotype as a risk factor for postmenopausal osteoporosis: *letter*, **636**
iron loading expression, Variation of iron loading expression in C282Y homozygous haemochromatosis probands and sib pairs: *letter*, **632**
iron storage, Classification and genetic features of neonatal haemochromatosis: a study of 27 affected pedigrees and molecular analysis of genes implicated in iron metabolism, **599**
isochromosome, Complex and segmental uniparental disomy (UPD): review and lessons from rare chromosomal complements: *review*, **497**
isochromosomes 2p and 2q, Uniparental isodisomy for paternal 2p and maternal 2q in a phenotypically normal female with two isochromosomes, i(2p) and i(2q): *letter*, **214**
isodisomy, Segmental uniparental isodisomy (UPD) for 2p16 without clinical symptoms: implications for UPD and other genetic studies of chromosome 2: *letter*, **106**

Jacobsen syndrome, Interstitial deletion of chromosome 11 (q22.3-q23.2) in a boy with mild developmental delay: *letter*, **621**
Japanese, Predisposing chromosome for spinocerebellar ataxia type 6 (SCA6) in Japanese: *letter*, **328**
jumping translocation, Cytogenetic and molecular study of a jumping translocation in a baby with Dandy-Walker malformation: *letter*, **67**

keratosis pilaris, Keratosis pilaris/ulerythema ophryogenes and 18p deletion: is it possible that the *LAMA1* gene is involved?: *letter*, **127**
kinship, Cultural aspects of cancer genetics: setting a research agenda: *review*, **425**
Kufor-Rakeb syndrome, Kufor-Rakeb syndrome, pallido-pyramidal degeneration with supranuclear upgaze paresis and dementia, maps to 1p36, **680**

laminin α2 chain, The expanding phenotype of laminin α2 chain (merosin) abnormalities: case series and review: *review*, **649**
laryngeal atresia, Laryngeal atresia, encephalocele, and limb deformities (LEL): a possible new syndrome: *letter*, **420**
***LBP-1c/CP2/LSF*,** Genetic association of an *LBP-1c/CP2/LSF* gene polymorphism with late onset Alzheimer's disease, **232**
Leigh syndrome, Mutations in *SURF1* are not specifically associated with Leigh syndrome: *letter*, **109**
leucodystrophy, Leucodystrophy and oculocutaneous albinism in a child with an 11q14 deletion, **35**
Li-Fraumeni syndrome, Detection of 11 germline inactivating *TP53* mutations and absence of *TP63* and *HCHK2* mutations in 17 French families with Li-Fraumeni or Li-Fraumeni-like syndrome: *letter*, **253**
Sensitivity and predictive value of criteria for *p53* germline mutation screening: *letter*, **43**
limb defects, Ocular malformations, postaxial polydactyly, and delayed intramembranous ossification: a new autosomal dominant condition: *letter*, **547**
linear skin defects, Molecular characterisation of a new case of microphthalmia with linear skin defects (MLS): *letter*, **411**
linkage analysis, A dominant gene for developmental dyslexia on chromosome 3, **658**
Development and application of linkage analysis in genetic diagnosis of familial hypertrophic cardiomyopathy: *letter*, **193**
Potential mapping of corneal dermoids to Xq24-qter: *letter*, **719**
liver, Association between the defective Pro369Ser mutation and in vivo intrahepatic α1-antitrypsin accumulation: *letter*, **472**
Classification and genetic features of neonatal haemochromatosis: a study of 27 affected pedigrees and molecular analysis of genes implicated in iron metabolism, **599**
long arm deletion, Mosaic terminal del(19)(q13.33:) in a girl with seizures and mental retardation: *letter*, **e2**
long QT syndrome, Mutation detection in long QT syndrome: a comprehensive set of primers and PCR conditions: *letter*, **705**
long term outcome, Acromicric dysplasia: long term outcome and evidence of autosomal dominant inheritance, **745**
loss of heterozygosity, Association of a novel constitutional translocation t(1q;3q) with familial renal cell carcinoma, **165**
lower incisors, Recessively inherited lower incisor hypodontia: *letter*, **551**
lymphoedema, Breakpoint analysis of Turner patients with partial Xp deletions: implications for the lymphoedema gene location, **591**
Clinical heterogeneity in lymphoedema-distichiasis with *FOXC2* truncating mutations, **761**

M34T, Genetic analysis of the connexin-26 M34T variant: identification of genotype M34T/M34T segregating with mild-moderate non-syndromic sensorineural hearing loss, **20**
Genetic analysis of the connexin-26 M34T variant: *letter*, **e24**
macrocephaly, Localisation of a gene for an autosomal recessive syndrome of macrocephaly, multiple epiphyseal dysplasia, and distinctive facies to chromosome 15q26, **369**
macrogonadism, A new case of exomphalos, short limbs, and macrogonadism syndrome: *letter*, **e8**
magnetic resonance angiography, Autosomal dominant polycystic kidney disease unlinked to the *PKD1* and *PKD2* loci presenting as familial cerebral aneurysm: *letter*, **238**
male breast cancer, Male breast cancer in Cowden syndrome patients with germline *PTEN* mutations, **159**
male infertility, Alternative centromeric inactivation in a pseudodicentric t(Y;13)(q12;p11.2) translocation chromosome associated with extreme oligozoospermia: *letter*, **802**
Mosaicism for 45,X cell line may accentuate the severity of spermatogenic defects in men with AZFc deletion: *letter*, **798**
mannose binding lectin, The mannose binding lectin gene influences the severity of chronic liver disease in cystic fibrosis, **310**
manuscript submission, *JMG* Online: exploiting the potential of electronic publication and manuscript submission: *editorial*, **809**
maternal inheritance, Maternal gene effect in neurofibromatosis 2: fact or artefact?: *letter*, **783**
Maternally inherited duplication of the possible imprinted 14q31 region: *letter*, **343**
maternal serum positive triple test screening, Pregnancy outcome and long term prognosis in 868 children born after second trimester amniocentesis for maternal serum positive triple test screening and normal prenatal karyotype: *letter*, **336**
maternal uniparental disomy 1, Supernumerary marker chromosome (1) of paternal origin and maternal uniparental disomy 1 in a developmentally delayed child : *letter*, **885**
maternal UPD(7), Do patients with maternal uniparental disomy for chromosome 7 have a distinct mild Silver-Russell phenotype?: *letter*, **273**
***MECP2* gene,** Rett syndrome and the *MECP2* gene: *review*, **217**

***MECP2* mutation,** Evidence of somatic mosaicism for a *MECP2* mutation in females with Rett syndrome: diagnostic implications : *letter*, **867**
MECP2 mutation in non-fatal, non-progressive encephalopathy in a male, **171**
Angelman syndrome phenotype associated with mutations in *MECP2*, a gene encoding a methyl CpG binding protein, **224**
***MECP2*,** Segregation of a totally skewed pattern of X chromosome inactivation in four familial cases of Rett syndrome without *MECP2* mutation: implications for the disease, **435**
medullary thyroid carcinoma, Familial medullary thyroid carcinoma and prominent corneal nerves associated with the germline V804M and V778I mutations on the same allele of *RET*: *letter*, **784**
medulloblastoma, A distinct splice form of *APC* is highly expressed in neurones but not commonly mutated in neuroepithelial tumours: *letter*, **257**
melatonin, β_1-adrenergic antagonists improve sleep and behavioural disturbances in a circadian disorder, Smith-Magenis syndrome, **586**
Menière, A common ancestor for *COCH* related cochleovestibular (DFNA9) patients in Belgium and The Netherlands bearing the P51S mutation: *letter*, **61**
Mennonite, A G339R mutation in the *CTNS* gene is a common cause of nephropathic cystinosis in the south western Ontario Amish Mennonite population: *letter*, **615**
mental retardation, Cryptic telomeric rearrangements in subjects with mental retardation associated with dysmorphism and congenital malformations: *letter*, **417**
How microsatellite analysis can be exploited for subtelomeric chromosomal rearrangement analysis in mental retardation: *letter*, **e1**
Mosaic terminal del(19)(q13.33:) in a girl with seizures and mental retardation: *letter*, **e2**
Use of a set of highly polymorphic minisatellite probes for the identification of cryptic 1p36.3 deletions in a large collection of patients with idiopathic mental retardation: *letter*, **121**
merosin, The expanding phenotype of laminin α2 chain (merosin) abnormalities: case series and review: *review*, **649**
metachromatic leucodystrophy, Multiple metachromatic leucodystrophy alleles in an unaffected subject: a case of dispermic chimaerism: *letter*, **e15**
methods, A comparison of methods for gene dosage analysis in HMSN type 1, **90**
methylation, Methylation matters: *review*, **285**
microdeletion, Microdeletion in the *FMR-1* gene: an apparent null allele using routine clinical PCR amplification: *letter*, **624**
microdeletion 2p, A case of ring chromosome 2 with growth retardation, mild dysmorphism, and microdeletion of 2p detected using FISH: *letter*, **e32**
microdissection, A supernumerary marker chromosome with a neocentromere derived from 5p14→pter: *letter*, **559**
microphthalmia, Molecular characterisation of a new case of microphthalmia with linear skin defects (MLS): *letter*, **411**
Ocular malformations, postaxial polydactyly, and delayed intramembranous ossification: a new autosomal dominant condition: *letter*, **547**
microsatellite analysis, How microsatellite analysis can be exploited for subtelomeric chromosomal rearrangement analysis in mental retardation: *letter*, **e1**
microsatellite instability, Hereditary and somatic DNA mismatch repair gene mutations in sporadic endometrial carcinoma: *letter*, **461**
mid frequency hearing loss, A cysteine substitution in the zona pellucida domain of α-tectorin results in autosomal dominant, postlingual, progressive, mid frequency hearing loss in a Spanish family: *letter*, **13**
migraine, Increased risk of sensorineural hearing loss and migraine in patients with a rare mitochondrial DNA variant 4336A>G in $tRNA^{Gln}$: *letter*, **400**
minisatellite, Use of a set of highly polymorphic minisatellite probes for the identification of cryptic 1p36.3 deletions in a large collection of patients with idiopathic mental retardation: *letter*, **121**
mitochondria, A frameshift mitochondrial complex I gene mutation in a patient with dystonia and cataracts: is the mutation pathogenic?: *letter*, **58**
mitochondrial DNA, A frameshift mitochondrial complex I gene mutation in a patient with dystonia and cataracts: is the mutation pathogenic?: *letter*, **58**
Mitochondrial DNA inheritance in patients with deleted mtDNA: *letter*, **28**
Prevalence of mitochondrial DNA mutations in childhood/congenital onset non-syndromal sensorineural hearing impairment, **229**
mitochondrial genome, Homoplasmic 3316G→A in the *ND1* gene of the mitochondrial genome: a pathogenic mutation or a neutral polymorphism?: *letter*, **e10**
mitochondrial $tRNA^{Tyr}$ mutation (5877G→A), Functional characterisation of mitochondrial $tRNA^{Tyr}$ mutation (5877G→A) associated with familial chronic progressive external ophthalmoplegia: *letter*, **703**
MMR, Screening families with endometrial and colorectal cancers for germline mutations: *letter*, **e29**
modulating gene, The mannose binding lectin gene influences the severity of chronic liver disease in cystic fibrosis, **310**
Moebius syndrome, Molecular cytogenetic characterisation of a complex 46,XY,t(7;8;11;13) chromosome rearrangement in a patient with Moebius syndrome: *letter*, **117**
molecular genetics, Molecular genetic heterogeneity in autosomal dominant drusen, **381**
molecular studies, Use of a set of highly polymorphic minisatellite probes for the identification of cryptic 1p36.3 deletions in a large collection of patients with idiopathic mental retardation: *letter*, **121**
molecular testing, Eight years' experience of direct molecular testing for myotonic dystrophy in Wales: *letter*, **e42**
Mondini malformation, Split hand/split foot malformation associated with sensorineural deafness, inner and middle ear malformation, hypodontia, congenital vertical talus, and deletion of eight microsatellite markers in 7q21.1-q21.3: *letter*, **405**
mortality, An epidemiological study of Wolf-Hirschhorn syndrome: life expectancy and cause of mortality, **674**
Anderson-Fabry disease: clinical manifestations and impact of disease in a cohort of 60 obligate carrier females: *letter*, **769**
Anderson-Fabry disease: clinical manifestations and impact of disease in a cohort of 98 hemizygous males, **750**
mosaicism, No evidence for mosaicism in Silver-Russell syndrome: *letter*, **e11**
MRC/CFAS study, Genetic association of an *LBP-1c/CP2/LSF* gene polymorphism with late onset Alzheimer's disease, **232**
mRNA, Hair roots as the ideal source of mRNA for genetic testing: *letter*, **e20**
***MRX49*,** Absence of learning difficulties in a hyperactive boy with a terminal Xp deletion encompassing the *MRX49* locus: *letter*, **466**
***MSH6*,** Atypical HNPCC owing to *MSH6* germline mutations: analysis of a large Dutch pedigree, **318**
mtDNA, Frequency of mitochondrial transfer RNA mutations and deletions in 225 patients presenting with respiratory chain deficiencies, **665**
Heterogeneous tissue distribution of a mitochondrial DNA polymorphism in heteroplasmic subjects without mitochondrial disorders, **312**
mtDNA variant 4336A>G, Increased risk of sensorineural hearing loss and migraine in patients with a rare mitochondrial DNA variant 4336A>G in $tRNA^{Gln}$: *letter*, **400**
multiple endocrine neoplasia type 1, Founder effect in multiple endocrine neoplasia type 1 (MEN 1) in Finland: *letter*, **185**
multiple epiphyseal dysplasia, Localisation of a gene for an autosomal recessive syndrome of macrocephaly, multiple epiphyseal dysplasia, and distinctive facies to chromosome 15q26, **369**
multiple isolated neurofibromas, A clinical study of patients with multiple isolated neurofibromas: *letter*, **485**
mUPD(7), No evidence for mosaicism in Silver-Russell syndrome: *letter*, **e11**
Silver-Russell syndrome: a dissection of the genetic aetiology and candidate chromosomal regions: *review*, **810**
mutation, Association between the defective Pro369Ser mutation and in vivo intrahepatic α1-antitrypsin accumulation: *letter*, **472**
De novo mutations in the 5' regulatory region of the Norrie disease gene in retinopathy of prematurity: *letter*, **e46**
Disruption of one intra-chain disulphide bond in the carboxyl-terminal propeptide of the proα1(I) chain of type I procollagen permits slow assembly and secretion of overmodified, but stable procollagen trimers and results in mild osteogenesis imperfecta, **443**
High frequency of the *ApoB*-100 R3500Q mutation in Bulgarian hypercholesterolaemic subjects: *letter*, **536**
Limited contribution of interchromosomal gene conversion to *NF1* gene mutation: *letter*, **481**
mutation analysis, GATA3 abnormalities and the phenotypic spectrum of HDR syndrome, **374**
Mutation detection in long QT syndrome: a comprehensive set of primers and PCR conditions: *letter*, **705**
mutation detection, Interpreting epidemiological research: blinded comparison of methods used to estimate the prevalence of inherited mutations in *BRCA1* , **824**
mutation Y279C, Mutations in the glycine receptor α1 subunit (*GLRA1*) gene in hereditary hyperekplexia pedigrees: evidence for non-penetrance of mutation Y279C: *letter*, **e17**
mutations, *BRCA1* and *BRCA2* mutation status and cancer family history of Danish women affected with multifocal or bilateral breast cancer at a young age, **361**
No evidence of allelic heterogeneity in the *DYT1* gene of European patients with early onset torsion dystonia: *letter*, **e35**
Novel mutations of *TMPRSS3* in four DFNB8/B10 families segregating congenital autosomal recessive deafness: *letter*, **396**
myosin light chain mutations, Myosin light chain mutations in familial hypertrophic cardiomyopathy: phenotypic presentation and frequency in Danish and South African populations: *letter*, **e43**
myotonic dystrophy, Eight years' experience of direct molecular testing for myotonic dystrophy in Wales: *letter*, **e42**
Molecular and clinical study of two myotonic dystrophy homozygotes: *letter*, **e40**

Presymptomatic testing in myotonic dystrophy: genetic counselling approaches : *letter*, **846**
Trinucleotide repeat contraction: a pitfall in prenatal diagnosis of myotonic dystrophy : *letter*, **850**

nail-patella syndrome, The small patella syndrome: description of five cases from three families and examination of possible allelism with familial patella aplasia-hypoplasia and nail-patella syndrome: *letter*, **209**
natural history, Anderson-Fabry disease: clinical manifestations and impact of disease in a cohort of 98 hemizygous males, **750**
***NBS1* gene,** Atypical clinical picture of the Nijmegen breakage syndrome associated with developmental abnormalities of the brain: *letter*, **e3**
NBS1, A novel mutation and novel features in Nijmegen breakage syndrome: *letter*, **113**
Nijmegen breakage syndrome in a Dutch patient not resulting from a defect in *NBS1*: *letter*, **e19**
***ND1* gene,** Homoplasmic 3316G→A in the *ND1* gene of the mitochondrial genome: a pathogenic mutation or a neutral polymorphism?: *letter*, **e10**
neocentromere, A supernumerary marker chromosome with a neocentromere derived from 5p14→pter: *letter*, **559**
neonatal, Classification and genetic features of neonatal haemochromatosis: a study of 27 affected pedigrees and molecular analysis of genes implicated in iron metabolism, **599**
neonatal hypertrypsinaemia, Analysis of the entire coding region of the cystic fibrosis transmembrane regulator gene in neonatal hypertrypsinaemia with normal sweat test: *letter*, **202**
nephropathic cystinosis, A G339R mutation in the *CTNS* gene is a common cause of nephropathic cystinosis in the south western Ontario Amish Mennonite population: *letter*, **615**
neural crest cells, *GDNF* as a candidate modifier in a type 1 neurofibromatosis (NF1) enteric phenotype, **638**
neurocristopathy, A heterozygous endothelin 3 mutation in Waardenburg-Hirschsprung disease: is there a dosage effect of *EDN3*/*EDNRB* gene mutations on neurocristopathy phenotypes?: *letter*, **205**
neurodevelopment, A majority of fragile X males with methylated, full mutation alleles have significant levels of *FMR1* messenger RNA, **453**
neurofibromas, A clinical study of patients with multiple isolated neurofibromas: *letter*, **485**
neurofibromatosis 2, Maternal gene effect in neurofibromatosis 2: fact or artefact?: *letter*, **783**
neurofibromatosis type 1 (NF1), Limited contribution of interchromosomal gene conversion to *NF1* gene mutation: *letter*, **481**
NF1, *GDNF* as a candidate modifier in a type 1 neurofibromatosis (NF1) enteric phenotype, **638**
Nijmegen breakage syndrome, A novel mutation and novel features in Nijmegen breakage syndrome: *letter*, **113**
Atypical clinical picture of the Nijmegen breakage syndrome associated with developmental abnormalities of the brain: *letter*, **e3**
Nijmegen breakage syndrome in a Dutch patient not resulting from a defect in *NBS1*: *letter*, **e19**
non-Robertsonian translocation, Stable non-Robertsonian dicentric chromosomes: four new cases and a review: *letter*, **76**
Norrie disease, De novo mutations in the 5' regulatory region of the Norrie disease gene in retinopathy of prematurity: *letter*, **e46**

obituary, Cyril Astley Clarke, **281**
Cyril Clarke, *Journal of Medical Genetics*, and the foundation of clinical genetics, **283**
Persisting memories of Cyril Clarke in Baltimore, **284**
obligate carriers, Anderson-Fabry disease: clinical manifestations and impact of disease in a cohort of 60 obligate carrier females: *letter*, **769**
oculocutaneous albinism, Leucodystrophy and oculocutaneous albinism in a child with an 11q14 deletion, **35**
oestrogens, *SHOX* haploinsufficiency and overdosage: impact of gonadal function status: *review*, **1**
oligodontia, Anauxetic dysplasia, a spondylometaepiphyseal dysplasia with extreme dwarfism: *letter*, **262**
oncology nurse training, Oncology nurse training in cancer genetics: *letter*, **691**
osteogenesis imperfecta, Disruption of one intra-chain disulphide bond in the carboxyl-terminal propeptide of the proα1(I) chain of type I procollagen permits slow assembly and secretion of overmodified, but stable procollagen trimers and results in mild osteogenesis imperfecta, **443**
Homozygosity for a splice site mutation of the *COL1A2* gene yields a non-functional proα2(I) chain and an EDS/OI clinical phenotype: *letter*, **132**
ovarian cancer, Surveillance or surgery? A description of the factors that influence high risk premenopausal women's decisions about prophylactic oophorectomy, **683**

p53, Sensitivity and predictive value of criteria for *p53* germline mutation screening: *letter*, **43**
pachydermoperiostosis, A syndrome of overgrowth and acromegaloidism with normal growth hormone secretion is associated with chromosome 11 pericentric inversion: *letter*, **338**
Papillon-Lefèvre syndrome, Evidence of a founder effect for four cathepsin C gene mutations in Papillon-Lefèvre syndrome patients, **96**
PAR1, Triplication of several PAR1 genes and part of the *Homo sapiens* specific Yp11.2/Xq21.3 region of homology in a 46,X,t(X;Y)(p22 33;p11.2) male with schizophrenia: *letter*, **710**
Parkinson's disease, Kufor-Rakeb syndrome, pallido-pyramidal degeneration with supranuclear upgaze paresis and dementia, maps to 1p36, **680**
partial deletion, A partial deletion of the aspartoacylase gene is the cause of Canavan disease in a family from Mexico: *letter*, **e9**
partial deletion of chromosome 3p, Interstitial deletion of 3p22.2-p24.2: the first reported case: *letter*, **349**
partial trisomy, Clinical and cytogenetic characterisation of a patient with Down syndrome resulting from a 21q22.1→qter duplication: *letter*, **73**
partial trisomy mosaicism, Maternal uniparental isodisomy 11q13→qter in a dysmorphic and mentally retarded female with partial trisomy mosaicism 11q13→qter : *letter*, **876**
patella aplasia-hypoplasia syndrome, The small patella syndrome: description of five cases from three families and examination of possible allelism with familial patella aplasia-hypoplasia and nail-patella syndrome: *letter*, **209**
Pearson syndrome, Mitochondrial DNA inheritance in patients with deleted mtDNA: *letter*, **e28**
peroxisome biogenesis disorders, Temperature sensitive acyl-CoA oxidase import in group A peroxisome biogenesis disorders: *letter*, **871**
Peters' anomaly, Phenotypic heterogeneity of *CYP1B1*: mutations in a patient with Peters' anomaly: *letter*, **324**
Peutz-Jeghers disease, Peutz-Jeghers families unlinked to *STK11*/*LKB1* gene mutations are highly predisposed to primitive biliary adenocarcinoma, **356**
phenotype-genotype correlation, Clinical and molecular characterisation of 80 patients with 5p deletion: genotype-phenotype correlation, **151**
phenotypic spectrum, GATA3 abnormalities and the phenotypic spectrum of HDR syndrome, **374**
phosphomannomutase, A broad spectrum of clinical presentations in congenital disorders of glycosylation I: a series of 26 cases, **14**
phosphomannose isomerase, A broad spectrum of clinical presentations in congenital disorders of glycosylation I: a series of 26 cases, **14**
physical mapping, Leucodystrophy and oculocutaneous albinism in a child with an 11q14 deletion, **35**
plasma fibrinogen, Fibrinogen genotypes (α and β) are associated with plasma fibrinogen levels in Chinese: *letter*, **e31**
point mutation, Prevalence of mitochondrial DNA mutations in childhood/congenital onset non-syndromal sensorineural hearing impairment, **229**
point mutations, *SHOX* point mutations in dyschondrosteosis: *letter*, **323**
polyendocrinopathy, Novel mutations of *FOXP3* in two Japanese patients with immune dysregulation, polyendocrinopathy, enteropathy, X linked syndrome (IPEX): *letter*, **874**
polymorphism, Heterogeneous tissue distribution of a mitochondrial DNA polymorphism in heteroplasmic subjects without mitochondrial disorders, **312**
The −48 CT polymorphism in the presenilin 1 promoter is associated with an increased risk of developing Alzheimer's disease and an increased Aβ load in brain, **353**
postmenopausal osteoporosis, Haptoglobin genotype as a risk factor for postmenopausal osteoporosis: *letter*, **636**
practical implications, *SHOX* haploinsufficiency and overdosage: impact of gonadal function status: *review*, **1**
Prader-Willi syndrome, Molecular characterisation of four cases of intrachromosomal triplication of chromosome 15q11-q14, **26**
Population prevalence and estimated birth incidence and mortality rate for people with Prader-Willi syndrome in one UK Health Region: *letter*, **792**
preaxial polydactyly, A novel acropectoral syndrome maps to chromosome 7q36, **304**
preconceptional carrier couple screening, Participation in preconceptional carrier couple screening: characteristics, attitudes, and knowledge of both partners: *letter*, **695**
predictive genetic testing, Comprehension of cancer risk one and 12 months after predictive genetic testing for hereditary non-polyposis colorectal cancer: *letter*, **787**
predictive testing, A community based study on intentions regarding predictive testing for hereditary breast cancer: *letter*, **540**
pre-eclampsia, A polymorphism in the gene for microsomal epoxide hydrolase is associated with pre-eclampsia, **234**
Thrombophilic polymorphisms in pre-eclampsia: altered frequency of the functional 98C>T polymorphism of glycoprotein IIIa: *letter*, **775**
pregnancy outcome, Pregnancy outcome and long term prognosis in 868 children born after second trimester amniocentesis for maternal serum positive triple test screening and normal prenatal karyotype: *letter*, **336**

premalignant laryngeal neoplasm, Role of *TP53* P72R polymorphism in human papillomavirus associated premalignant laryngeal neoplasm: *letter,* **327**
premolars, Recessively inherited lower incisor hypodontia: *letter,* **551**
prenatal diagnosis, Trinucleotide repeat contraction: a pitfall in prenatal diagnosis of myotonic dystrophy : *letter,* **850**
prenatal testing, Prenatal testing for Huntington's disease: experience within the UK 1994-1998: *letter,* **333**
presenilin, The −48 CT polymorphism in the presenilin 1 promoter is associated with an increased risk of developing Alzheimer's disease and an increased Aβ load in brain, **353**
presymptomatic testing, Presymptomatic testing in myotonic dystrophy: genetic counselling approaches : *letter,* **846**
prevalence, Anderson-Fabry disease: clinical manifestations and impact of disease in a cohort of 98 hemizygous males, **750**
Population prevalence and estimated birth incidence and mortality rate for people with Prader-Willi syndrome in one UK Health Region: *letter,* **792**
***PRF1* gene,** Six novel mutations in the *PRF1* gene in children with haemophagocytic lymphohistiocytosis: *letter,* **643**
primary care, Recurrent mutations in the deafness gene *GJB2* (connexin 26) in British Asian families: *letter,* **530**
Pro369Ser, Association between the defective Pro369Ser mutation and in vivo intrahepatic α1-antitrypsin accumulation: *letter,* **472**
procollagen, Disruption of one intra-chain disulphide bond in the carboxyl-terminal propeptide of the proα1(I) chain of type I procollagen permits slow assembly and secretion of overmodified, but stable procollagen trimers and results in mild osteogenesis imperfecta, **443**
prognathism, Anauxetic dysplasia, a spondylometaepiphyseal dysplasia with extreme dwarfism: *letter,* **262**
promoter, The −48 CT polymorphism in the presenilin 1 promoter is associated with an increased risk of developing Alzheimer's disease and an increased Aβ load in brain, **353**
prophylactic oophorectomy, Surveillance or surgery? A description of the factors that influence high risk premenopausal women's decisions about prophylactic oophorectomy, **683**
Proteus syndrome, *PTEN* mutations are uncommon in Proteus syndrome: *letter,* **480**
proximal deletion 18q, Molecular characterisation of a proximal chromosome 18q deletion: *letter,* **128**
pseudogene, A unique form of autosomal dominant cataract explained by gene conversion between β-crystallin B2 and its pseudogene: *letter,* **392**
Limited contribution of interchromosomal gene conversion to *NF1* gene mutation: *letter,* **481**
Pseudoxanthoma elasticum: evidence for the existence of a pseudogene highly homologous to the *ABCC6* gene: *letter,* **457**
pseudoxanthoma elasticum, Pseudoxanthoma elasticum: evidence for the existence of a pseudogene highly homologous to the *ABCC6* gene: *letter,* **457**
psoriasis, Identification of a novel psoriasis susceptibility locus at 1p and evidence of epistasis between *PSORS1* and candidate loci, **7**
***PSORS1*,** Identification of a novel psoriasis susceptibility locus at 1p and evidence of epistasis between *PSORS1* and candidate loci, **7**
psychological studies, Psychological studies in Huntington's disease: making up the balance: *letter,* **852**
***PTEN* mutation,** The spectrum and evolution of phenotypic findings in *PTEN* mutation positive cases of Bannayan-Riley-Ruvalcaba syndrome: *letter,* **52**
***PTEN* mutations,** *PTEN* mutations are uncommon in Proteus syndrome: *letter,* **480**
***PTEN*,** A novel germline mutation of the *PTEN* gene in a patient with macrocephaly, ventricular dilatation, and features of VATER association, **820**
Male breast cancer in Cowden syndrome patients with germline *PTEN* mutations, **159**

quantitative PCR, Submicroscopic deletions of the *APC* gene: a frequent cause of familial adenomatous polyposis that may be overlooked by conventional mutation scanning: *letter,* **129**
quantitative RT-PCR, A majority of fragile X males with methylated, full mutation alleles have significant levels of *FMR1* messenger RNA, **453**

R3500Q, High frequency of the *ApoB*-100 R3500Q mutation in Bulgarian hypercholesterolaemic subjects: *letter,* **536**
reading disability, A dominant gene for developmental dyslexia on chromosome 3, **658**
reduced penetrance alleles, New problems in testing for Huntington's disease: the issue of intermediate and reduced penetrance alleles: *letter,* **e12**
respiratory chain deficiency, Frequency of mitochondrial transfer RNA mutations and deletions in 225 patients presenting with respiratory chain deficiencies, **665**
retinal dystrophy, Clustering and frequency of mutations in the retinal guanylate cyclase (*GUCY2D*) gene in patients with dominant cone-rod dystrophies: *letter,* **611**
Does Bardet-Biedl syndrome have a characteristic face?: *letter,* **e14**
retinitis pigmentosa, Evaluation of the *ELOVL4* gene in families with retinitis pigmentosa linked to the *RP25* locus: *letter,* **478**
retinopathy of prematurity, De novo mutations in the 5' regulatory region of the Norrie disease gene in retinopathy of prematurity: *letter,* **e46**
Rett syndrome, Angelman syndrome phenotype associated with mutations in *MECP2*, a gene encoding a methyl CpG binding protein, **224**
Evidence of somatic mosaicism for a *MECP2* mutation in females with Rett syndrome: diagnostic implications : *letter,* **867**
MECP2 mutation in non-fatal, non-progressive encephalopathy in a male, **171**
Rett syndrome and the *MECP2* gene: *review,* **217**
Segregation of a totally skewed pattern of X chromosome inactivation in four familial cases of Rett syndrome without *MECP2* mutation: implications for the disease, **435**
***RET*,** Familial medullary thyroid carcinoma and prominent corneal nerves associated with the germline V804M and V778I mutations on the same allele of *RET*: *letter,* **784**
ring chromosome 2, A case of ring chromosome 2 with growth retardation, mild dysmorphism, and microdeletion of 2p detected using FISH: *letter,* **e32**
risk estimation, Peutz-Jeghers families unlinked to *STK11/LKB1* gene mutations are highly predisposed to primitive biliary adenocarcinoma, **356**
risk factor, Haptoglobin genotype as a risk factor for postmenopausal osteoporosis: *letter,* **636**
risk perception, Risk perception and cancer worry: an exploratory study of the impact of genetic risk counselling in women with a family history of breast cancer: *letter,* **139**
RNA splicing, Homozygosity for a splice site mutation of the *COL1A2* gene yields a non-functional proα2(I) chain and an EDS/OI clinical phenotype: *letter,* **132**
Roberts syndrome, A case of Roberts syndrome described in 1737: *letter,* **565**
Robertsonian translocation, Complex and segmental uniparental disomy (UPD): review and lessons from rare chromosomal complements: *review,* **497**
***RP25*,** Evaluation of the *ELOVL4* gene in families with retinitis pigmentosa linked to the *RP25* locus: *letter,* **478**

S288K allele, Identification of a transcriptionally compromised allele of *c-MYC* in a North American family: *letter,* **47**
Saethre-Chotzen syndrome, Pure partial 7p trisomy including the *TWIST*, *HOXA*, and *GLI3* genes: *letter,* **178**
satellited short arms, Satellites on the terminal short arm of chromosome 12 (12ps), inherited through several generations in three families: a new variant without phenotypic effect: *letter,* **723**
SC phocomelia, A case of Roberts syndrome described in 1737: *letter,* **565**
SCA6, Twenty CAG repeats are sufficient to cause the SCA6 phenotype: *letter,* **e38**
schizophrenia, Triplication of several PAR1 genes and part of the *Homo sapiens* specific Yp11.2/Xq21.3 region of homology in a 46,X,t(X;Y)(p22 33;p11.2) male with schizophrenia: *letter,* **710**
screening, Science or black magic?: *letter,* **e16**
The importance of good images in using hair to screen for breast cancer: *letter,* **e16**
***SEDL*,** A missense mutation in the *SEDL* gene results in delayed onset of X linked spondyloepiphyseal dysplasia in a large pedigree: *letter,* **409**
segmental disomy, Segmental uniparental isodisomy (UPD) for 2p16 without clinical symptoms: implications for UPD and other genetic studies of chromosome 2: *letter,* **106**
segmental hypopigmentation, A novel mutation in the endothelin B receptor gene in a patient with Shah-Waardenburg syndrome and Down syndrome: *letter,* **646**
segmental uniparental disomy, Maternal uniparental isodisomy 11q13→qter in a dysmorphic and mentally retarded female with partial trisomy mosaicism 11q13→qter : *letter,* **876**
seizures, Mosaic terminal del(19)(q13.33:) in a girl with seizures and mental retardation: *letter,* **e2**
sensorineural deafness, Interstitial deletion of 3p22.2-p24.2: the first reported case: *letter,* **349**
Split hand/split foot malformation associated with sensorineural deafness, inner and middle ear malformation, hypodontia, congenital vertical talus, and deletion of eight microsatellite markers in 7q21.1-q21.3: *letter,* **405**
sensorineural hearing loss, Increased risk of sensorineural hearing loss and migraine in patients with a rare mitochondrial DNA variant 4336A>G in $tRNA^{Gln}$: *letter,* **400**
Shah-Waardenburg syndrome, A novel mutation in the endothelin B receptor gene in a patient with Shah-Waardenburg syndrome and Down syndrome: *letter,* **646**
SHFM, Split hand/split foot malformation associated with sensorineural deafness, inner and middle ear malformation, hypodontia, congenital vertical talus, and deletion of eight microsatellite markers in 7q21.1-q21.3: *letter,* **405**
short limbs, A new case of exomphalos, short limbs, and macrogonadism syndrome: *letter,* **e8**
short stature, Interstitial deletion of 3p22.2-p24.2: the first reported case: *letter,* **349**

***SHOX*, *SHOX* haploinsufficiency and overdosage: impact of gonadal function status: *review*, 1**
SHOX point mutations in dyschondrosteosis: *letter*, 323
Silver-Russell phenotype, Do patients with maternal uniparental disomy for chromosome 7 have a distinct mild Silver-Russell phenotype?: *letter*, 273
Silver-Russell syndrome, Identification of interstitial maternal uniparental disomy (UPD) (14) and complete maternal UPD(20) in a cohort of growth retarded patients, 86
No evidence for mosaicism in Silver-Russell syndrome: *letter*, e11
Silver-Russell syndrome: a dissection of the genetic aetiology and candidate chromosomal regions: *review*, 810
Sir Cyril Clarke, Cyril Astley Clarke, 281
Cyril Clarke, *Journal of Medical Genetics*, and the foundation of clinical genetics, 283
Persisting memories of Cyril Clarke in Baltimore, 284
situs inversus, Suggestive linkage of situs inversus and other left-right axis anomalies to chromosome 6p: *letter*, 182
skeletal muscle, The expanding phenotype of laminin α2 chain (merosin) abnormalities: case series and review: *review*, 649
skewed X chromosome inactivation, Segregation of a totally skewed pattern of X chromosome inactivation in four familial cases of Rett syndrome without *MECP2* mutation: implications for the disease, 435
small patella syndrome, The small patella syndrome: description of five cases from three families and examination of possible allelism with familial patella aplasia-hypoplasia and nail-patella syndrome: *letter*, 209
Smith-Magenis syndrome, β_1-adrenergic antagonists improve sleep and behavioural disturbances in a circadian disorder, Smith-Magenis syndrome, 586
***SMN1* deletions,** Detection of heterozygous *SMN1* deletions in SMA families using a simple fluorescent multiplex PCR method: *letter*, 240
***SOX10* mutations,** Novel mutations of *SOX10* suggest a dominant negative role in Waardenburg-Shah syndrome: *letter*, e30
spastic paraplegia, The spastic paraplegia SPG10 locus: narrowing of critical region and exclusion of sodium channel gene *SCN8A* as a candidate: *letter*, 65
spectral karyotyping (SKY), Molecular cytogenetic characterisation of a complex 46,XY,t(7;8;11;13) chromosome rearrangement in a patient with Moebius syndrome: *letter*, 117
spermatogenesis, Mosaicism for 45,X cell line may accentuate the severity of spermatogenic defects in men with AZFc deletion: *letter*, 798
SPG10 locus, The spastic paraplegia SPG10 locus: narrowing of critical region and exclusion of sodium channel gene *SCN8A* as a candidate: *letter*, 65
SPG3 locus, Hereditary spastic paraplegia linked to chromosome 14q11-q21: reduction of the SPG3 locus interval from 5.3 to 2.7 cM: *letter*, e39
spinal muscular atrophy, Detection of heterozygous *SMN1* deletions in SMA families using a simple fluorescent multiplex PCR method: *letter*, 240
***SPINK1* gene mutations,** Low prevalence of *SPINK1* gene mutations in adult patients with chronic idiopathic pancreatitis: *letter*, 243
spinocerebellar ataxia, Twenty CAG repeats are sufficient to cause the SCA6 phenotype: *letter*, 1
spinocerebellar ataxia type 6, Predisposing chromosome for spinocerebellar ataxia type 6 (SCA6) in Japanese: *letter*, 328
splice form, A distinct splice form of *APC* is highly expressed in neurones but not commonly mutated in neuroepithelial tumours: *letter*, 257
splicing mutation, A novel 3600+11.5 kb C>G homozygous splicing mutation in a black African, consanguineous CF family: *letter*, e4
Cystic fibrosis patients with the 3272-26A>G splicing mutation have milder disease than F508del homozygotes: a large European study: *letter*, 777
sponastrime dysplasia, Sponastrime dysplasia: presentation in infancy : *letter*, 889
spondyloepiphyseal dysplasia, A missense mutation in the *SEDL* gene results in delayed onset of X linked spondyloepiphyseal dysplasia in a large pedigree: *letter*, 409
***SRY* gene,** Analysis of the *SRY* gene in Turner syndrome patients with Y chromosomal material: *letter*, e41
stable dicentric chromosomes, Stable non-Robertsonian dicentric chromosomes: four new cases and a review: *letter*, 76
sternal deformity, A novel acropectoral syndrome maps to chromosome 7q36, 304
***STGD3*,** Evaluation of the *ELOVL4* gene in families with retinitis pigmentosa linked to the *RP25* locus: *letter*, 478
subarachnoid haemorrhage, Autosomal dominant polycystic kidney disease unlinked to the *PKD1* and *PKD2* loci presenting as familial cerebral aneurysm: *letter*, 238
submicroscopic, Submicroscopic subtelomeric 1qter deletions: a recognisable phenotype?: *letter*, 175
submicroscopic subtelomeric rearrangements, Clinical studies on submicroscopic subtelomeric rearrangements: a checklist, 145
subtelomere deletions, Submicroscopic subtelomeric 1qter deletions: a recognisable phenotype?: *letter*, 175
subtelomeric chromosomal rearrangement, How microsatellite analysis can be exploited for subtelomeric chromosomal rearrangement analysis in mental retardation: *letter*, e1
sulphate transporter gene, Sulphate transporter gene mutations in apparently isolated club foot: *letter*, 191
supernumerary marker chromosome, A supernumerary marker chromosome with a neocentromere derived from 5p14→pter: *letter*, 559
supernumerary marker chromosome 1, Supernumerary marker chromosome (1) of paternal origin and maternal uniparental disomy 1 in a developmentally delayed child : *letter*, 885
supernumerary ring chromosome, Molecular characterisation of a supernumerary ring chromosome in a patient with VATER association: *letter*, e6
***SURF1*,** Mutations in *SURF1* are not specifically associated with Leigh syndrome: *letter*, 109

TBX1, Mutation analysis of *TBX1* in non-deleted patients with features of DGS/VCFS or isolated cardiovascular defects, e45
***TBX5* deletion,** Detection of a large *TBX5* deletion in a family with Holt-Oram syndrome, e44
terminal deletion 15q, De novo terminal deletion of chromosome 15q26.1 characterised by comparative genomic hybridisation and FISH with locus specific probes: *letter*, 617
tetrasomy distal 5p, A supernumerary marker chromosome with a neocentromere derived from 5p14→pter: *letter*, 559
thrombophilic polymorphisms, Thrombophilic polymorphisms in pre-eclampsia: altered frequency of the functional 98C>T polymorphism of glycoprotein IIIa: *letter*, 775
tibial/radial aplasia, Laryngeal atresia, encephalocele, and limb deformities (LEL): a possible new syndrome: *letter*, 420
***TMPRSS3*,** Novel mutations of *TMPRSS3* in four DFNB8/B10 families segregating congenital autosomal recessive deafness: *letter*, 396
torsion dystonia, No evidence of allelic heterogeneity in the *DYT1* gene of European patients with early onset torsion dystonia: *letter*, e35
***TP53* mutations,** Detection of 11 germline inactivating *TP53* mutations and absence of *TP63* and *HCHK2* mutations in 17 French families with Li-Fraumeni or Li-Fraumeni-like syndrome: *letter*, 253
TP53 P72R polymorphism, Role of *TP*53 P72R polymorphism in human papillomavirus associated premalignant laryngeal neoplasm: *letter*, 327
transforming growth factor β receptor type II, Germline mutation analysis of the transforming growth factor β receptor type II (*TGFBR2*) and E-cadherin (*CDH1*) genes in early onset and familial colorectal cancer: *letter*, e7
translocation, Association of a novel constitutional translocation t(1q;3q) with familial renal cell carcinoma, 165
translocation t(4;8), Two further cases of WHS with unbalanced de novo translocation t(4;8) characterised by CGH and FISH: *letter*, e21
trinucleotide repeat, A majority of fragile X males with methylated, full mutation alleles have significant levels of *FMR1* messenger RNA, 453
Trinucleotide repeat contraction: a pitfall in prenatal diagnosis of myotonic dystrophy : *letter*, 850
tRNA mutations, Frequency of mitochondrial transfer RNA mutations and deletions in 225 patients presenting with respiratory chain deficiencies, 665
$tRNA^{Gln}$, Increased risk of sensorineural hearing loss and migraine in patients with a rare mitochondrial DNA variant 4336A>G in $tRNA^{Gln}$: *letter*, 400
***TSC2*,** A novel missense mutation in the GTPase activating protein homology region of *TSC2* in two large families with tuberous sclerosis complex: *letter*, 347
tuberous sclerosis complex, A novel missense mutation in the GTPase activating protein homology region of *TSC2* in two large families with tuberous sclerosis complex: *letter*, 347
tumour suppressor, Two hits revisited again: *review*, 81
Turner syndrome, Analysis of the *SRY* gene in Turner syndrome patients with Y chromosomal material: *letter*, e41
Breakpoint analysis of Turner patients with partial Xp deletions: implications for the lymphoedema gene location, 591
***TWIST* mutations,** Identification of a new *TWIST* mutation (7p21) with variable eyelid manifestations supports locus homogeneity of BPES at 3q22: *letter*, 470
***TWIST*,** Pure partial 7p trisomy including the *TWIST*, *HOXA*, and *GLI3* genes: *letter*, 178
two hit model, Two hits revisited again: *review*, 81

***UGT1A1*,** Interaction of coding region mutations and the Gilbert-type promoter abnormality of the *UGT1A1* gene causes moderate degrees of unconjugated hyperbilirubinaemia and may lead to neonatal kernicterus: *letter*, 244
ulerythema ophryogenes, Keratosis pilaris/ulerythema ophryogenes and 18p deletion: is it possible that the *LAMA1* gene is involved?: *letter*, 127
uniparental disomy, Ellis-van Creveld syndrome resulting from segmental uniparental disomy of chromosome 4: *letter*, e18

Segmental uniparental isodisomy (UPD) for 2p16 without clinical symptoms: implications for UPD and other genetic studies of chromosome 2: *letter*, **106**

uniparental disomy 14, Identification of interstitial maternal uniparental disomy (UPD) (14) and complete maternal UPD(20) in a cohort of growth retarded patients, **86**

uniparental disomy 20, Identification of interstitial maternal uniparental disomy (UPD) (14) and complete maternal UPD(20) in a cohort of growth retarded patients, **86**

uniparental disomy (UPD), Complex and segmental uniparental disomy (UPD): review and lessons from rare chromosomal complements: *review*, **497**

uniparental isodisomy, Uniparental isodisomy for paternal 2p and maternal 2q in a phenotypically normal female with two isochromosomes, i(2p) and i(2q): *letter*, **214**

United Kingdom Study, Prenatal testing for Huntington's disease: experience within the UK 1994-1998: *letter*, **333**

urogenital malformations, B cell immunodeficiency, distal limb abnormalities, and urogenital malformations in a three generation family: a novel autosomal dominant syndrome?: *letter*, **488**

van der Woude syndrome, Genetic heterogeneity and exclusion of a modifying locus at 17p11.2-p11.1 in Finnish families with van der Woude syndrome: *letter*, **198**

VATER, A novel germline mutation of the *PTEN* gene in a patient with macrocephaly, ventricular dilatation, and features of VATER association, **820**

VATER association, Molecular characterisation of a supernumerary ring chromosome in a patient with VATER association: *letter*, **e5**

velocardiofacial syndrome, Mutation analysis of *TBX1* in non-deleted patients with features of DGS/VCFS or isolated cardiovascular defects, **e45**

vestibular, A common ancestor for *COCH* related cochleovestibular (DFNA9) patients in Belgium and The Netherlands bearing the P51S mutation: *letter*, **61**

VHL morbidity, *VHL* c.505 T>C mutation confers a high age related penetrance but no increased overall mortality, **508**

VHL mortality, *VHL* c.505 T>C mutation confers a high age related penetrance but no increased overall mortality, **508**

***VHL* gene,** *VHL* c.505 T>C mutation confers a high age related penetrance but no increased overall mortality, **508**

villous atrophy, Mitochondrial DNA inheritance in patients with deleted mtDNA: *letter*, **e28**

von Hippel-Lindau disease, Association of a novel constitutional translocation t(1q;3q) with familial renal cell carcinoma, **165**

Waardenburg-Hirschsprung disease, A heterozygous endothelin 3 mutation in Waardenburg-Hirschsprung disease: is there a dosage effect of *EDN3*/*EDNRB* gene mutations on neurocristopathy phenotypes?: *letter*, **205**

Waardenburg-Shah syndrome, Novel mutations of *SOX10* suggest a dominant negative role in Waardenburg-Shah syndrome: *letter*, **e30**

***WFS1*,** Wolfram syndrome: a clinical and molecular genetic analysis: *letter*, **e37**

Williams syndrome, Coeliac disease in Williams syndrome, **767**

Wolf-Hirschhorn syndrome, An epidemiological study of Wolf-Hirschhorn syndrome: life expectancy and cause of mortality, **674**

Two further cases of WHS with unbalanced de novo translocation t(4;8) characterised by CGH and FISH: *letter*, **e21**

Wolfram syndrome, Wolfram syndrome: a clinical and molecular genetic analysis: *letter*, **e37**

X chromosome, Segregation of a totally skewed pattern of X chromosome inactivation in four familial cases of Rett syndrome without *MECP2* mutation: implications for the disease, **435**

X inactivation, The androgen receptor and DXS15-134 markers show a high rate of discordance for germline X chromosome inactivation: *letter*, **474**

X linked, Potential mapping of corneal dermoids to Xq24-qter: *letter*, **719**

X linked inheritance, A missense mutation in the *SEDL* gene results in delayed onset of X linked spondyloepiphyseal dysplasia in a large pedigree: *letter*, **409**

Xp11.4, Breakpoint analysis of Turner patients with partial Xp deletions: implications for the lymphoedema gene location, **591**

Xp22.3, Molecular characterisation of a new case of microphthalmia with linear skin defects (MLS): *letter*, **411**

Xq21.3/Yp11.2, Triplication of several PAR1 genes and part of the *Homo sapiens* specific Yp11.2/Xq21.3 region of homology in a 46,X,t(X;Y)(p22 33;p11.2) male with schizophrenia: *letter*, **710**

XX male, Triplication of several PAR1 genes and part of the *Homo sapiens* specific Yp11.2/Xq21.3 region of homology in a 46,X,t(X;Y)(p22 33;p11.2) male with schizophrenia: *letter*, **710**

Y chromosome, Analysis of the *SRY* gene in Turner syndrome patients with Y chromosomal material: *letter*, **e41**

Mosaicism for 45,X cell line may accentuate the severity of spermatogenic defects in men with AZFc deletion: *letter*, **798**

Y;autosome translocation, Alternative centromeric inactivation in a pseudodicentric t(Y;13)(q12;p11.2) translocation chromosome associated with extreme oligozoospermia: *letter*, **802**

zona pellucida, A cysteine substitution in the zona pellucida domain of α-tectorin results in autosomal dominant, postlingual, progressive, mid frequency hearing loss in a Spanish family: *letter*, **e13**